

KiActiv HealthCheck Bibliography

Abbott, R.D., White, L.R., Ross, G.W., Masaki, K.H., Curb, J.D., and Petrovitch, H. (2004). Walking and dementia in physically capable elderly men. *Journal of the American Medical Association*, **292**, 1447-1453.

Academy Health Inequalities Forum (2015). *Exercise: The miracle cure and the role of the doctor in promoting it*. Academy of Medical Royal Colleges: London, UK.

Adams-Campbell, L.L., Rosenberg, L., Rao, R.S., and Palmer, J.R. (2001). Strenuous physical activity and breast cancer risk in African-American women. *Journal of the National Medical Association*, **93(7-8)**, 267-275.

Ahlskog, J.E., Geda, Y.E., Graff-Radford, N.R., and Peterson, R.C. (2011). Physical exercise as a preventative or disease modifying treatment of dementia and brain aging. *Mayo Clinic Proceedings*, **86(9)**, 876-884.

Ainsworth, B.E., Haskell, W.L., Whitt, M.C., Irwin, M.L., Swartz, A.M., Strath, S.J., O'Brien, W.L., Bassett, D.R., Schmitz, K.H., Emplaincourt, P.O., Jacobs, D.R., and Leon, A.S. (2000). Compendium of physical activities: an update of activity codes and MET intensities. *Medicine and Science in Sports and Exercise*, **32(Supplement 9)**, S498-S504.

Ainsworth, B.E., Leon, A.S., Richardson, M.T., Jacobs, D.R., and Paffenbarger, R.S. (1993). Accuracy of the College Alumni Physical Activity Questionnaire. *Journal of Clinical Epidemiology*, **46**, 1403-1411.

American College of Sports Medicine (2005). *ACSM Guidelines for Exercise Testing and Prescription*, Seventh Edition. Baltimore: Lippincott Williams and Wilkins.

American College of Sports Medicine (1998). *ACSM's Resource Manual for Guidelines for Exercise Testing and Prescription*, Third edition. Baltimore: Lippincott Williams and Wilkins.

American College of Sports Medicine (1990). The recommended quality and quantity of exercise for developing and maintaining fitness in healthy adults. *Medicine and Science in Sports and Exercise*, **22**, 265-274.

Andersen, R.E., Wadden, T.A., Bartlett, S.J., Zemel, B., Verde, T.J., and Franckowiak, S.C. (1999). Effects of lifestyle activity vs. Structured aerobic exercise in obese women. *Journal of the American Medical Association*, **281**, 335-340.

Arem, H., Moore, S.C., Patel, A., Hartge, P., Berrington de Gonzalez, A., Visvanathan, K., Campbell, P.T., Freedman, M., Weiderpass, E., Adami, H.O., Linet, M.S., Lee, I-M., and Matthews, C.E. (2015). Leisure time physical activity and mortality: A detailed pooled analysis of the dose-response relationship. *JAMA Internal Medicine*, doi:10.1001/jamainternmed.2015.0533.

Aresu, M., Bécares, L., Brage, S., Chaudhury, M., Doyle-Francis, M., Eslinger, D., Fuller, E., Gunning, N., Hall, J., Hirani, V., Jotangia, D., Mindell, J., Moody, A., Ogunbadejo, T., Pickup, D., Reilly, N.,

Robinson, C., Roth, M., and Wardle, H. (2009). *Health survey for England 2008 volume 1. Physical activity and fitness*. UK: The NHS Information Centre for Health and Social Care.

Bak, E.E., Hellénus, M-L. and Ekblom, B. (2010). Are we facing a new paradigm of inactivity physiology? *British Journal of Sports Medicine*, doi:10.1136/bjism.2009.067702

Bang, H., Edwards, A.M., Bombback, A.S., Ballantyne, C.M., Brillon, D., Callahan, M.A., Teutsch, S.M., Mushlin, A.I., and Kern, L.M. (2009). A patient self-assessment diabetes screening score: development, validation, and comparison to other diabetes risk assessment scores. *Annals of Internal Medicine*, **151(11)**, 775-783.

Barker, D.J.P., Eriksson, J.G., Forsén, T., and Osmond, C. (2002). Fetal origins of adult disease: Strength of effects and biological basis. *International Journal of Epidemiology*, **31(6)**, 1235-1239.

Barlow, C.E., LaMonte, M.J., FitzGerald, S.J., Kampert, J.B., Perrin, J.L., and Blair, S.N. (2006). Cardiorespiratory fitness is an independent predictor of hypertension incidence among initially normotensive healthy women. *American Journal of Epidemiology*, **163(2)**, 142-150.

Barnes, D.E. and Yaffe, K. (2011). The projected effect of risk factor reduction on Alzheimer's disease prevalence. *Lancet: Neurology*, doi: 10.1016/S1474-4422(11)70072-2.

Barulli, D. and Stern, Y. (2013) Efficiency, capacity, compensation, maintenance, plasticity: emerging concepts in cognitive reserve. *Trends in Cognitive Sciences*, **17**, 502-509.

Bassuk, S.S. and Manson, J.E. (2005). Epidemiological evidence for the role of physical activity in reducing risk of type 2 diabetes and cardiovascular disease. *Journal of Applied Physiology*, **99**, 1193-1204.

Batty, D. and Thune, I. (2000). Does physical activity prevent cancer? Evidence suggests protection against colon and probably breast cancer. *British Medical Journal*, **321**, 1424-1425.

Bauman, A. (2012). Don't just sit there – Stand up. Sydney School of Public Health, Sydney Medical School.

Bauman, W.A. and Spungen, A.M. (2008). Coronary heart disease in individuals with spinal cord injury: assessment of risk factors. *Spinal Cord*, **46(7)**, 466-476.

Belsky, D.W., Caspi, A., Houts, R., Cohen, H.J., Corcoran, D.L., Danese, A., Harrington, H., Israel, S., Schaefer, J.D., Sugden, K., Williams, B., Yashin, A.I., Poulton, R., and Moffitt, T.E. (2015). Quantification of biological aging in young adults. *Proceedings of the National Academy of Sciences of the United States of America*, Published ahead of print on 6th July 2015, doi:10.1073/pnas.1506264112.

Ben-Ezra, V., Jankowski, C., Kendrick, K., and Nichols, D. (1995). Effect of intensity and energy expenditure on postexercise insulin responses in women. *Journal of Applied Physiology*, **79**, 2029-2034.

Benedict, C., Brooks, S.J., Kullberg, J., Nordenskjöld, R., Burgos, J., Le Grevès, M., Kilander, L., Larsson, E.M., Johansson, L., Ahlström, H., Lind, L., and Schiöth, H.B. (2012). Association between physical activity and brain health in older adults. *Neurobiology of Aging*, **34(1)**, 83-90.

Berlin, J. and Colditz, G.A. (1990). A meta-analysis of physical activity in the prevention of coronary heart disease. *American Journal of Epidemiology*, **132**, 612-628.

Bernstein, L., Patel, A.V., Ursin, G., Sullivan-Halley, J., Press, M.F., Deapen, D., Berlin, J.A., Daling, J.R., McDonald, J.A., Norman, S.A., Malone, K.E., Strom, B.L., Liff, J., Folger, S.G., Simon, M.S., Burkman, R.T., Marchbanks, P.A., Weiss, L.K., and Spirtas, R. (2005). Lifetime recreational exercise activity and breast cancer risk among black women and white women. *Journal of the National Cancer Institute*, **97(22)**, 1671-1679.

Bernstein, L., Henderson, B.E., Hanisch, R., Sullivan-Halley, J., and Ross, R.K. (1994). Physical exercise and reduced risk of breast cancer in young women. *Journal of the National Cancer Institute*, **86**, 1403-1408.

Bey, L. and Hamilton, M.T. (2003). Suppression of skeletal muscle lipoprotein lipase activity during physical inactivity: a molecular reason to maintain daily low-intensity activity. *Journal of Physiology*, **551(Pt 2)**, 673-682.

Beydoun, M.A., Beydoun, H.A., Gamaldo, A.A., Teel, A., Zonderma, A.B., and Wang, Y. (2014). Epidemiologic studies of modifiable factors associated with cognition and dementia: systematic review and meta-analysis. *BMC Public Health*, **14**, 643 doi:10.1186/1471-2458-14-643.

Bishop J.G. and S.G. Aldana (1999). *Step Up to Wellness*. Boston: Allyn & Bacon Publishing.

Biswas, A., Oh, P.I., Faulkner, G.E., Bajaj, R.R., Silver, M.A., Mitchell, M.S., and Alter, D.A. (2015). Sedentary time and its association with risk for disease incidence, mortality, and hospitalisation in adults. *Annals of Internal Medicine*, **162(2)**, 123-132.

Black, A., Coward, W., Cole, T., and Prentice, A. (1996). Human energy expenditure in affluent societies: an analysis of 574 doubly labelled water measurements. *European Journal of Clinical Nutrition*, **50**, 72-92.

Blair, S.N. (2009) Physical Inactivity: the biggest public health problem of the 21st century. *British Journal of Sports Medicine*, **43(1)**, 1-2.

Blair, S.N., Kohl, H.W. and Barlow, C.E. (1993) Physical activity, physical fitness, and all-cause mortality in women: do women need to be active? *Journal of the American College of Nutrition*, **12**, 368-371.

Blanck, H.M., McCullough, M.L., Patel, A.V., Gillespie, C., Calle, E.E., Cokkinides, V.E., Galuska, D.A., Khan, L.K., and Serdula, M.K. (2007). Sedentary behaviour, recreational physical activity, and 7-year weight gain among postmenopausal U.S. women. *Obesity*, **15(6)**, 1578-1588.

Block, J.E., Friedlander, A.L., Brooks, G.A., Steiger, P., Stubbs, H.A., and Genant, H.K. (1989). Determinants of bone density among athletes engaged in weight-bearing and non-weight-bearing activity. *Journal of Applied Physiology*, **67**, 110-1105.

Block, J.E., Genant, H.K. and Black, D. (1986). Greater vertebral bone mineral mass in exercising young men. *The Western Journal of Medicine*, **145**, 39-42.

Blondell, S.J., Hammersley-Mather, R. and Veerman, J.L. (2014). Does physical activity prevent cognitive decline and dementia?: a systematic review and meta-analysis of longitudinal studies. *BMC Public Health*, **14**, 510 doi:10.1186/1471-2458-14-510.

Bogardus, C., Lillioja, S., Ravussin, E., Abbott, W., Zawadzki, J.K., Young, A., Knowler, W.C., Jacobowitz, R., and Moll, P.P. (1986). Familial dependence of the resting metabolic rate. *New England Journal of Medicine*, **315**, 96-100.

Borer, K.T. (2005). Physical activity in the prevention and amelioration of osteoporosis in women: interaction of mechanical, hormonal and dietary factors. *Sports Medicine*, **35(9)**, 779-830.

Bouchard, C. and Shephard, R.J. (1994). Physical activity, fitness and health: the model and key concepts. In: *Physical Activity, Fitness, and Health*. Bouchard, C., Shephard, R.J. and Stephens, T. (Eds.) Champaign, IL: Human Kinetics, 1994, pp.77-88.

Bouchard, C., Shephard, R.J. and Stephens, T. (Eds.; 1994). Physical activity, fitness, and health. International proceedings and consensus statement. Champaign, IL: Human Kinetics, 1994.

Brahm, H., Ström, H., Piehl-Aulin, K., Mallmin, H., and Ljunghall, S. (1997). Bone metabolism in endurance trained athletes: a comparison to population-based controls based on DXA, SXA, quantitative ultrasound, and biochemical markers. *Calcified Tissue International*, **61**, 448-454.

Braith, R.W., Pollock, M.L., Lowenthal, D.T., Graves, J.E., and Limacher, M.C. (1994). Moderate- and high-intensity exercise lowers blood pressure in normotensive subjects 60 to 79 years of age. *The American Journal of Cardiology*, **73(15)**, 1124-1128.

British Medical Association (2012). *Healthy transport, healthy lives*. BMA: London, UK.

Brodsky, H., Heffernan, M., Kochan, N.A., Draper, B., Trollor, J.N., Reppermund, S., Slavin, M.J., and Sachdev, P.S. (2013). Mild cognitive impairment in a community sample: the Sydney Memory and Ageing Study.

Brooks, J.H.M. and Ferro, A. (2012). The physician's role in prescribing physical activity for the prevention and treatment of essential hypertension. *JRSM Cardiovascular Disease*, **1(4)**, 12.

Brooks, G.A., Butte, N.F., Rand, W.M., Flatt, J.P., and Caballero, B. (2004). Chronicle of the Institute of Medicine physical activity recommendations: how a physical activity recommendation came to be among dietary recommendations. *American Journal of Clinical Nutrition*, **79**, 921-930.

Brown, W.J., Bauman, A.E., Bull, F.C., and Burton, N.W. (2012). *Development of Evidence-based Physical Activity Recommendations for Adults (18-64 years)*. Report prepared for the Australian Government Department of Health, August 2012.

Brown, W.J., Burton, N.W. and Rowan, P.J. (2007). Updating the evidence on physical activity and health in women. *American Journal of Preventive Medicine*, **33(5)**, 404-411/

Brown, W.J., Hockey, R., and Dobson, A.J. (2011). Physical activity, sitting and weight gain in Australian women. *Journal of Science and Medicine in Sport*, **14(7)**, S93

Brown, W.J., McLaughlin, D., Leung, J., McCaul, K.A., Flicker, L., Almeida, O.P., Hankey, G.J., Lopez, D., and Dobson, A.J. (2012). Physical activity and all-cause mortality in older women and men. *British Journal of Sports Medicine*, **46**, 664-668.

Brown, W., Williams, L., Ford, J., Ball, K., and Dobson, A. (2005). Identifying the 'energy gap': magnitude and determinants of five year weight gain in mid age women. *Obesity Research*, **13(8)**, 1431-1441.

Bull, F., Biddle, S., Buchner, D., Ferguson, R., Foster, C., Fox, K., Haskell, B., Mutrie, N., Murphy, M., Reilly, J., Riddoch, C., Skelton, D., Stratton, G., Tremblay, M., and Watts, C. (2010). Physical activity guidelines in the UK: Review and recommendations. School of Sport, Exercise and Health Sciences, Loughborough University, May 2010.

Buchman, A.S., Boyle, P.A., Wilson, R.S., Bienias, J.L., and Bennett, D.A. (2007). Physical activity and motor decline in older persons. *Muscle and Nerve*, **35**, 354-362.

Buchman, A.S., Leurgans, S.E., Boyle, P.A., Schneider, J.A., Arnold, S.E., and Bennett, D.A. (2011). Combinations of motor measures more strongly predict adverse health outcomes in old age: the Rush Memory and Aging Project, a community-based cohort study. *BMC Medicine*, **9**, 42.

Buchman, A.S., Yu, L., Boyle, P.A., Shah, R.C., Bennett, D.A. (2012). Total daily physical activity and longevity in old age. *Archives of Internal Medicine*, **172**, 444-446.

Buman, M.P., Heckler, E.B., Haskell, W.L., Pruitt, L., Conway, T.L., Cain, K.L., Sallis, J.F., Saelens, B.E., Frank, L.D., and King, A.C. (2010). Objective light-intensity physical activity associations with rated health in older adults. *American Journal of Epidemiology*, **172(10)**, 1155-1165.

Buman, M.P., Winkler, E.A., Kurka, J.M., Heler, E.B., Baldwin, C.M., Owen, N., Answorth, B.E., Healy, G.N., and Gardiner, P.A. (2014). Reallocating time to sleep, sedentary behaviours, or active behaviours: associations with cardiovascular disease risk biomarkers, NHANES 2005-2006. *American Journal of Epidemiology*, **179(3)**, 323-334.

Burgomaster, K.A., Howarth, K.R., Phillips, S., Rakobowchuk, M., MacDonald, M.J., McGee, S.L., and Gibala, M.J. (2008). Similar metabolic adaptations during exercise after low volume sprint interval and traditional endurance training in humans. *Journal of Physiology*, **586**, 151-160.

Byberg, L., Melhus, H., Gedeberg, R., Sundstrom, J., Ahlbom, A., Zethelius, B., Berglund, L.G., Wolk, A., and Michaëlsson, K. (2009) Total mortality after changes in leisure time physical activity in 50 year old men: 35 year follow-up of population based cohort. *British Medical Journal*, **338**, b688.

Calbet, J.A., Moysi, J.S., Dorado, C., and Rodriguez, L.P. (1998). Bone mineral content and density in professional tennis players. *Calcified Tissue International*, **62**, 491-496.

Carpenter, C.L., Ross, R.K., Paganini-Hill, A., and Bernstein, L. (1999). Lifetime exercise activity and breast cancer risk among post-menopausal women. *British Journal of Cancer*, **80**, 1852-1858.

Carson, V., Ridgers, N.D., Howard, B.J., Winkler, E.A.H., Healy, G.N., Owen, N., Dunstan, D.W., and Salmon, J. (2013). Light-intensity physical activity and cardiometabolic biomarkers in US adolescents. *PLoS ONE*, **8**(8), e71417.

Caspersen, C.J., Powell, K.E. and Christenson, G.M. (1985). Physical activity, exercise, and physical fitness: definitions and distinctions for health-related research. *Public Health Reports*, **100**(2), 126-131.

Centers for Disease Control and Prevention (CDC; 2010). National health and nutrition examination survey data 2003-2004, 2005-2006. Available at :
http://www.cdc.gov/nchs/nhanes/nhanes_questionnaires.htm.

Centers for Disease Control and Prevention (2008). Physical Activity for Everyone – How much physical activity do adults need?
<http://www.cdc.gov/physicalactivity/everyone/guidelines/adults.html>.

Chakravarthy, M.V. and Booth, F.W. (2004). Eating, exercise, and “thrifty” genotypes: connecting the dots towards an evolutionary understanding of modern chronic diseases. *Journal of Applied Physiology*, **96**, 3-10.

Chao, A., Connell, C.J., Jacobs, E.J., McCullough, M.L., Patel, A.V., Calle, E.E., Cokkinides, V.E., and Thun, M.J. (2004). Amount, type, and timing of recreational physical activity in relation to colon and rectal cancer in older adults: the Cancer Prevention Study II Nutrition Cohort. *Cancer Epidemiology, Biomarkers and Prevention*, **13**(12), 2187-2195.

Chau, J.Y., Grunseit, A.C., Chey, T., Stamatakis, E., Brown, W.J., Matthews, C.E., Bauman, A.E., and van der Ploeg, H.P. (2013). Daily sitting time and all-cause mortality: A meta-analysis. *PLoS ONE*, **8**(11): e80000. doi:10.1371/journal.pone.0080000.

Chen, L., Magliano, D.J., Balkau, B., Colagiuri, S., Zimmet, P.Z., Tonkin, A.M., Mitchell, P., Phillips, P.J., and Shaw, J.E. (2010). AUSDRISK: an Australian Type 2 Diabetes Risk Assessment Tool based on demographic, lifestyle and simple anthropometric measures. *Medical Journal of Australia*, **192**(4), 197-202.

Chief Medical Officer (2004). At least five a week: Evidence on the impact of physical activity and its relationship to health – A report from the Chief Medical Officer. Department of Health, http://webarchive.nationalarchives.gov.uk/+dh.gov.uk/en/publicationsandstatistics/publications/publicationspolicyandguidance/dh_4080994

Chief Medical Officer (2009). Annual report 2009, Department of Health. http://www.sthc.co.uk/Documents/CMO_Report_2009.pdf

Chief Medical Officers of England, Scotland, Wales and Northern Ireland (2011). Start active, stay active: a report on physical activity from the four home countries' Chief Medical Officers, Department of Health, <https://www.gov.uk/government/publications/start-active-stay-active-a-report-on-physical-activity-from-the-four-home-countries-chief-medical-officers>

Chobanian, A.V., Bakris, G.L., Black, H.R., Cushman, W.C., Green, L.A., Izzo, J.L., Jones, D.W. Materson, B.J., Oparil, S., Wright, J.T., and Roccella, E.J. (2003). The seventh report of the joint national

committee on prevention, detection, evaluation, and treatment of high blood pressure: the JNC 7 report. *Journal of the American Medical Association*, **289(19)**, 2560-2572.

Chomistek, A.K., Manson, J.E., Stefanick, M.L., Lu, B., Sands-Lincoln, M., Going, S.B., Garcia, L., Allison, M.A., Sims, S.T., LaMonte, M.J., Johnson, K.C., and Eaton, C.B. (2013). The relationship of sedentary behaviour and physical activity to incident cardiovascular disease: Results from the women's health initiative. *Journal of the American College of Cardiology*, **61(23)**, 2346-2354.

Cleroux, J., Feldman, R.D. and Petrella, R.J. (1999). Recommendations on physical exercise training. *Canadian Medical Association Journal*, **160**, S21-S28.

Colcombe, S. and Kramer, A.F. (2003). Fitness effects on the cognitive function of older adults: a meta-analytic study. *Psychological Science*, **14(2)**, 125-130.

Colcombe, S.J., Erickson, K.I., Scaf, P.E., Kim, J.S., Prakash, R., McAuley, E., Elavsky, S., Marquez, D.X., Hu, L., and Kramer, A.F. (2006). Aerobic exercise training increases brain volume in aging humans. *Journal of Gerontology: Series A, Biological Sciences and Medical Sciences*, **61(11)**, 1166-1170.

Colletti, L.A., Edwards, J., Gordon, L., Shary, J., and Bell, N.H. (1989). The effects of muscle-building exercise on bone mineral density of the radius, spine, and hip in young men. *Classified Tissue International*, **45**, 12-14.

Colley, R.C., Garrigué, D., Janssen, I., Craig, C.L., Clarke, J., and Tremblay, M.S. (2011). Physical activity of Canadian adults: accelerometer results from the 2007 to 2009 Canadian Health Measures Survey. *Health Reports*, **22(1)**, 7-14.

Cooper, D.M., Barstow, T.J., Bergner, A., and Lee, W.N. (1989). Blood glucose turnover during high- and low-intensity exercise. *American Journal of Physiology*, **257**, E405-E412.

Cordain, L., Latin, R.W. and Behnke, J.J. (1986). The effects of an aerobic running program on bowel transit time. *Journal of Sports Medicine and Physical Fitness*, **26**, 101-104.

Cornelissen, V.A. and Fagard, R.H. (2005). Effects of endurance training on blood pressure, blood pressure-regulating mechanisms, and cardiovascular risk factors. *Hypertension*, **46**, 667-675.

Cox, K.L., Puddey, I.B., Burke, V., Beilin, L.J., Morton, A.R., and Bettridge, H.F. (1996). Determinants of change in blood pressure during S.W.E.A.T.: the sedentary women exercise adherence trial. *Clinical and Experimental Pharmacology and Physiology*, **23**, 567-569.

Crouse, S.F., O'Brien, B.C., Grandjean, P.W., Lowe, R.C., Rohack, J.J., Green, J.S., and Tolson, H. (1997). Training intensity, blood lipids, and apolipoproteins in men with high cholesterol. *Journal of Applied Physiology*, **82(1)**, 270-277.

Danese, A., and McEwan, B.S. (2012). Adverse childhood experiences, allostasis, allostatic load, and age-related disease. *Physiology and Behaviour*, **106(1)**, 29-39.

Davies, E.J., Moxham, T., Rees, K., Singh, S., Coats, A.J., Ebrahim, S., Lough, F., and Taylor, R.S. (2010). Exercise based rehabilitation for heart failure. *Cochrane Database of Systematic Reviews*, **14(4)**, CD003331.

Department of Health (2011a). *Reducing obesity and improving diet, 11th July 2011*. Department of Health: London, UK.

Department of Health (2011). Health Survey for England – 2010, Trend tables. Publication date: 15th December, 2011. <http://www.hscic.gov.uk/pubs/hse10trends>

Department of Health (2009). Be Active, Be Healthy: a plan to get the nation moving. <http://webarchive.nationalarchives.gov.uk/20090410133341/http://diabetes.nhs.uk/news-folder/be-active-be-healthy-a-plan-for-getting-the-nation-moving>

Department of Health (2004) *At least five a week; Evidence on the impact of physical activity and its relationship to health*. London: DoH.

Diabetes Prevention Program [DPP] Research Group (2003). Within-trial cost-effectiveness of lifestyle intervention or metformin for the primary prevention of type 2 diabetes. *Diabetes Care*, **26(9)**, 2518-2523.

Diabetes Prevention Program [DPP] Research Group (2002). The Diabetes Prevention program (DPP): Description of lifestyle intervention. *Diabetes Care*, **25(12)**, 2165-2171.

Dipietro, L. (1999). Physical activity in the prevention of obesity: current evidence and research issues. *Medicine and Science in Sports and Exercise*, **31**, S542-S546.

Donnelly, J.E., Blair, S.N., Jakicic, J.M., Manore, M.M., Rankin, J.W., and Smith, B.K. (2009). American College of Sports Medicine position stand. Appropriate physical activity intervention strategies for weight loss and prevention of weight regain for adults. *Medicine and Science in Sports and Exercise*. **41(2)**, 459–471.

Drygas W., Kostka, T., Jegier, A., and Kuński, H. (2000). Long term effects of different physical activity levels on coronary heart disease risk factors in middle-aged men. *International Journal of Sports Medicine*, **21(4)**, 235-41.

Drygas, W., Jegler, A. and Kunski, H. (1988) Study on threshold dose of physical activity in coronary heart disease prevention, part I: relationship between leisure time physical activity and coronary risk factors. *International Journal of Sports Medicine*, **9**, 275-278.

Duncan, J.J., Gordon, N.F., and Scott, C.B. (1991). Women walking for health and fitness: how much is enough? *Journal of the American Medical Association*, **266**, 3295-3299.

Dunn, A.L., Trivedi, M.H., Kampert, J.B., Clark, C.G., and Chambliss, H.O. (2005). Exercise treatment for depression: efficacy and dose response. *American Journal of Preventive Medicine*, **28(1)**, 1-8.

Dunn, A.L., Marcus, B.H., Kampert, J.B., Garcia, M.E., Kohl, H.W., and Blair, S.M. (1999). Comparison of lifestyle and structured interventions to increase physical activity and cardiorespiratory fitness: a randomised trial. *Journal of the American Medical Association*, **281**,327-334.

Dunstan, D.W., Kingwell, B.A., Larsen, R., Healy, G.N., Cerin, E., Hamilton, M.T., Shaw, J.E., Bertovic, D.A., Zimmet, P.Z., Salmon, J., and Owen, N. (2012). Breaking up prolonged sitting reduces postprandial glucose and insulin responses. *Diabetes Care*, **35(5)**, 96-983.

Dunstan, D.W., Barr, E.L.M., Healy, G.N., Salmon, J., Shaw, J.E., Balkau, B., Magliano, D.J., Cameron, A.J., Zimmet, P.Z., and Owen, N. (2010). Television viewing time and mortality: the AusDiab study. *Circulation*, **121**, 384-391.

Duppe, H., Gardsell, P., Johnell, O., Nilsson, B.E., and Ringsberg, K. (1997). Bone mineral density, muscle strength and physical activity. A population-based study of 332 subjects aged 15-42 years. *Acta Physiologica Scandinavica*, **68**, 97-103.

Durstine, J.L., Grandjean, P.W., Cox, C.A., and Thompson, P.D. (2002). Lipids, lipoproteins, and exercise. *Journal of Cardiopulmonary Rehabilitation*, **22(6)**, 385-398.

Durstine J.L. and W.L. Haskell (1994). Effects of exercise training on plasma lipids and lipoproteins. *Exercise and Sports Science Reviews*, **22**, 477-522.

Duvivier, B.M.F.M., Schaper, N.C., Bremers, M.A., van Crombrugge, G., Menheere, P.P.C.A., Kars, M., and Savelberg, H.H.C.M. (2013) Longer duration improves insulin action and plasma lipids more than shorter periods of moderate to vigorous exercise (cycling) in sedentary subjects when energy expenditure is comparable. *PLoS ONE*, **9(8)**, e105135. doi:10.1371/journal.pone.0105135.

Ekblom-Bak, E., Olsson, G., Ekblom, Ö., Ekblom, B., Bergström, G., and Börjesson, M. (2015). The daily movement pattern and fulfilment of physical activity recommendations in Swedish middle-aged adults: The SCAPIS pilot study. *PLoS ONE*, **10(5)**, e0126336. Doi:10.1371/journal.pone.0126336.

Elbaz, A., Vicente-Vytopilova, P., Tavernier, B., Sabia, S., Dumurgier, J., Mazoyer, B., Singh-Manoux, A., and Tzourio, C. (2013). Motor function in the elderly: evidence for the reserve hypothesis. *Neurology*, **81(5)**, 417-426.

Elley, C.R., Garrett, S., Rose, S.B., O'Dea, D., Lawton, B.A., Moyes, S.A., and Dowell, A.C. (2010). Cost-effectiveness of exercise on prescription with telephone support among women in general practice over 2 years. *British Journal of Sports Medicine*, doi:10.1136/bjism.2010.072439.

Erickson, K.I., Hillman, C.H. and Kramer, A.F. (2015). Physical activity, brain, and cognition. *Current Opinion in Behavioural Science*, **4**, 27-32.

Erickson, K., Gildengers, A.G. and Butters, M.A. (2013). Physical activity and brain plasticity in late adulthood. *Dialogues Clinical Neuroscience*, **15(1)**, 99-108.

Erickson, K.I., Voss, M.W., Prakash, R.S., Basak, C., Szabo, A., Chaddock, L., Kim, J.S., Heo, S., Alves, H., White, S.M., Wojcicki, T.R., Mailey, E., Vieira, V.J., Martin, S.A., Pence, B.D., Woods, J.A., McAuley, E., and Kramer, A.F. (2011). Exercise training increases size of hippocampus and improves memory. *PNAS*, **108(7)**, 3017-3022.

Erickson, K.I., Raji, C.A., Lopez, O.L., Becker, J.T., Rosano, C., Newman, A.B., Gach, H.M., Thompson, P.M., Ho, A.J., and Kuller, L.H. (2010). Physical activity predicts gray matter volume in late afternoon. *Neurology*, **75(16)**, 1415-1422.

Eriksson, J., Linström, J., Valle, T., Aunola, S., Härmäläinen, H., Ilanne-Parikka, P., Keinänen-Kiukaanniemi, S., Laakso, M., Lauhkonen, M., Lehto, P., Lehtonen, A., Louheranta, A., Mannelin, M., Martikkala, V., Rastas, M., Sundvall, J., Turpinen, A., Viljanen, T., Uusitupa, M., and Tuomilehto, J.

(1999). Prevention of type II diabetes in subjects with impaired glucose tolerance: the Diabetes Prevention Study (DPS) in Finland. Study design and 1-year interim report on the feasibility of the lifestyle intervention programme. *Diabetologia*, **42(7)**, 793-801.

Erlichman, J., Kerbey, A. and James, P. (2001). *Are current guidelines adequate to prevent unhealthy weight gain? A scientific appraisal for consideration by an Expert Panel of the International Obesity Task Force (IOTF)*. London, IOTF, pp113.

Erlichman, J., Kerbey, A. and James, P. (no date). Physical activity and its impact on health outcomes, Paper II: prevention of unhealthy weight gain and obesity by physical activity: An analysis of the evidence.

Fagard, R.H. (1999). Physical activity in the prevention and treatment of hypertension in the obese. *Medicine and Science in Sports and Exercise*, **31(11 Suppl)**, S624-S630.

Fagard, R.H. (2001). Exercise characteristics and the blood pressure response to dynamic physical training. *Medicine and Science in Sports and Exercise*, **33(6 Suppl)**, S484-S492.

Fagard, R.H. and Cornelissen, V.A. (2007). Effect of exercise on blood pressure control in hypertensive patients. *European Journal of Cardiovascular Prevention and Rehabilitation*, **14(1)**, 12-17.

Ferris, L.T., Williams, J.S., and Shen, C.L. (2007). The effect of acute exercise on serum brain-derived neurotrophic factor levels and cognitive function. *Medicine and Science in Sports and Exercise*, **39**, 728-734.

Ferro-Luzzi, A. and Martino, L. (1996). Obesity and physical activity. In *The origins and consequences of obesity*, pp. 207-227. Wiley Chichester, Ciba Foundation Symposium 201.

Fleischman, D.A., Yang, J., Arfanakis, K., Arvanitakis, Z., Leurgans, S.E., Turner, A.D., Barnes, L.L., Bennett, D.A., and Buchman, A.S. (2015). Physical activity, motor function, and white matter hyperintensity burden in healthy older adults. *Neurology*, **84**, 1-7.

Fletcher, G.F., Balady, G., Blair, S.N., Blumenthal, J., Caspersen, C., Chaitman, B., Epstein, S., Sivarajan Froelicher, E.S., Froelicher, V.F., Pina, I.L., and Pollock, M.L. (1996) Statement on Exercise: Benefits and Recommendations for Physical Activity Programs for All Americans. A Statement for Health Professionals by the Committee on Exercise and Cardiac Rehabilitation of the Council on Clinical Cardiology, American Heart Association. *Circulation*, **94**, 857-862.

Folsom, A.R., Kushi, L.H., and Hong, C.P. (2000). Physical activity and incident diabetes mellitus in postmenopausal women. *American Journal of Public Health*, **90**, 134-138.

Folsom, A.R., Prineas, R.J., Kaye, S.A., and Munger, R.G. (1990). Incidence of hypertension and stroke in relation to body fat distribution and other risk factors in older women. *Stroke*, **21**, 701-706.

Fong, D.Y.T., Ho, J.W.C., Hui, B.P.H., Lee, A.M., Macfarlane, D.J., Leung, S.S.K., Cerin, E., Chan, W.Y.Y., Leung, I.P.F., Lam, S.H.S., Taylor, A.J., and Cheng, K. (2012) Physical activity for cancer survivors: meta-analysis of randomised trials. *British Medical Journal*, **344**, e70.

Fontana, L., Kennedy, B.K., Longo, V.D., Seals, D., and Melov, S. (2014). Medical research: Are we ready? *Aging Cell*, 10.1111/ace.12338.

Food and Agriculture Organisation of the United Nations, World Health Organisation, and United Nations University (FAO/WHO/UNU; 2004). Human Energy Requirements, Report of a joint FAO/WHO/UNU Expert Consultation. Rome, 17-24 October 2001.

Fraser, G.E. and Shavlik, D. (1997). Risk factors, lifetime risk, and age at onset of breast cancer. *Annals of Epidemiology*, **7**, 375-382.

Friedenreich, C.M. (2011). Physical activity and breast cancer: review of the epidemiologic evidence and biologic mechanisms. *Recent Results in Cancer Research*, **188**, 125-139.

Friedenreich, C.M., Bryant, H.E., and Courneya, K.S. (2001). Case-control study of lifetime physical activity and breast cancer risk. *American Journal of Epidemiology*, **154(4)**, 336-347.

Friedlander, A.L., Casazza, G.A., Horning, M.A., Huie, M.J., Piacentini, M.F., Trimmer, J.K., and Brooks, G.A. (1998). Training-induced alterations of carbohydrate metabolism in women: women respond differently from men. *Journal of Applied Physiology*, **85**, 1175-1186.

Furberg, A.S. and Thune, I. (2003). Metabolic abnormalities (hypertension, hyperglycemia and overweight), lifestyle (high energy intake and physical inactivity) and endometrial cancer in a Norwegian cohort. *International Journal of Cancer*, **104**, 669-676.

Garcia-Aymerich, J., Lange, P., Benet, M., Schnohr, P., and Anto, J.M. (2006) Regular physical activity reduces hospital admission and mortality in chronic obstructive pulmonary disease: a population based cohort study. *Thorax*, **61**, 772-778.

Gavrilov, L.A., and Gavrilova, N.S. (2004). Early-life programming of aging and longevity: The idea of high initial damage load (the HDL hypothesis). *Annals of the New York Academy of Sciences*, **1019**, 496-501.

George, S.M., Alfano, C.M., Wilder Smith, A., Irwin, M.L., McTiernan, A., Bernstein, L., Baumgartner, K.B., and Ballard-Barbash, R. (2013). Sedentary behaviour, health-related quality of life and fatigue among breast cancer survivors. *Journal of Physical Activity and Health*, **10(3)**, 350-358.

Gillum, R.F., Mussolino, M.E. and Ingram, D.D. (1996). Physical activity and stroke incidence in women and men. The NHAMES I Epidemiologic Follow-up Study. *American Journal of Epidemiology*, **143(9)**, 860-869.

Gillman, M.W. (2005). Developmental origins of health and disease. *New England Journal of Medicine*, **353(17)**, 1848-1850.

Gibala, M.J. (2007). High-intensity interval training: a time-efficient strategy for health promotion? *Current Sports Medicine Reports*, **6**, 211-213.

Giovannucci, E., Ascherio, A., Rimm, E.B., Colditz, G.A., Stampfer, M.J., and Willett, W.C. (1995). Physical activity, obesity, and risk for colon cancer and adenoma in men. *Annals of Internal Medicine*, **122(5)**, 327-334.

Glümer, C., Carstensen, B., Sandbæk, A., Lauritzen, T., Jørgensen, T., and Borch-Johnsen, K. (2004). A Danish Diabetes Risk Score for Targeted Screening. *Diabetes Care*, **27(3)**, 727-733.

Goldstein, L.B. (2010). Physical activity and the risk of stroke. *Expert Review of Neurotherapeutics*, **10(8)**, 1263-1265.

Gons, R.A.R., Tuladhar, A.M., de Laat, K.F., van Vorden, A.G.W., van Dijk, E.J., and Norris, D.G. (2013). Physical activity is related to the structural integrity of cerebral white matter. *Neurology*, **81(11)**, 971-976.

Gordon, D.J., Probstfield, J.L., Garrison, R.J., Neaton, J.D., Castelli, W.P., Knoke, J.D., Jacobs, D.R.Jr., Bangdiwala, S., and Tyroler, H.A. (1989). High-density lipoprotein cholesterol and cardiovascular disease. Four prospective American studies. *Circulation*, **79(1)**, 8-15.

Gow, A.J., Bastin, M.E., Muñoz Maniega, S., Valdés Hernández, M.C., Morris, Z., Murray, C., Royle, N.A., Starr, J.M., Deary, I.J., Wardlaw, J.M. (2012). Neuroprotective lifestyles and the aging brain: activity, atrophy, and white matter integrity. *Neurology*, **79(17)**, 1802-1808.

Grandes, G., Sanchez, A., Sanchez-Pinilla, R.O., Torcal, J., Montoya, I., Lizarraga, K., and Serra, J. (2009). Effectiveness of physical activity advice and prescription by physicians in routine primary care: a cluster randomised trial. *Archives of Internal Medicine*, **169(7)**, 694-701.

Haapanen, N., Miilunpalo, S., Vuori, I., Oja, P., and Pasanen, M. (1997). Association of leisure time physical activity with the risk of coronary heart disease, hypertension and diabetes in middle-aged men and women. *International Journal of Epidemiology*, **26**, 739-747.

Hagberg, J.M., Montain, S.J., Martin, W.H., and Ehsani, M.A. (1989). Effect of exercise training in 60-69-year-old persons with essential hypertension. *American Journal of Cardiology*, **64**, 348-353.

Halbert, J.A., Silacy, C.A., Finucane, P., Withers, R.T., Hamdorf, P.A., and Andrews, G.R. (1997). The effectiveness of exercise training in lowering blood pressure: a meta-analysis of randomised controlled trials of 4 weeks or longer. *Journal of Human Hypertension*, **11**, 641-649.

Hambrecht, R., Niebauer, J., Marburger, C., Grunze, M., Kälberer, B., Hauer, K., Schlierf, G., Kübler, W., and Schuler, G. (1993). Various intensities of leisure time physical activity in patients with coronary artery disease: effects on cardiorespiratory fitness and progression of coronary atherosclerotic lesions. *Journal of the American College of Cardiology*, **22(2)**, 468-477.

Hamburg, N.M., McMackin, C.J., Huang, A.L., Shenouda, S.M., Widlansky, M.E., Schulz, E., Gokce, N., Ruderman, N.B., Keaney, J.F.Jr., and Vita, J.A. (2007). Physical inactivity rapidly induces insulin resistance and microvascular dysfunction in healthy volunteers. *Arteriosclerosis, Thrombosis and Vascular Biology*, **27(12)**, 2650-2656.

Hamer, M. and Chida, Y. (2009). Physical activity and risk of neurodegenerative disease: a systematic review of prospective evidence. *Psychological Medicine*, **39**, 3-11.

Hamer, M., Ingle, L., Carroll, S., and Stamatakis, E. (2012). Physical activity and cardiovascular mortality risk: possible protective mechanisms? *Medicine and Science in Sports and Exercise*, **44(1)**, 84-88.

Hamilton, M.T., Hamilton, D.G. and Zderic, T.W. (2007). Role of low energy expenditure and sitting on obesity, metabolic syndrome, type 2 diabetes and cardiovascular disease. *Diabetes*, **56**, 2655-2667.

Hamilton, M.T., Hamilton, D.G. and Zderic, T.W. (2004). Exercise physiology versus inactivity physiology: an essential concept for understanding lipoprotein lipase regulation. *Exercise and Sport Science Reviews*, **32(4)**, 161-166.

Hamman, R.F., Wing, R.R., Edelstein, S.L., Lachin, J.M., Bray, G.A., Delahanty, L., Hoskin, M., Kriska, A.M., Mayer-Davis, E.J., Pi-Sunyer, X., Regensteiner, J., Venditti, B., and Wylie-Rosett, J. (2006). Effect of weight loss with lifestyle intervention on risk of diabetes. *Diabetes Care*, **29(9)**, 2102-2107.

Hansen, A.S., Carstensen, B., Helge, J.W., Johansen, N.B., Gram, B., Christiansen, J.S., Brage, S., Lauritzen, T., Jørgensen, M.E., Aadahl, M., and Witte, D.R. (2013). Combined heart rate- and accelerometer-assessed physical activity energy expenditure and associations with glucose homeostasis markers in a population at high risk of developing diabetes: the ADDITION-PRO study. *Diabetes Care*, **36**, 3062-3069.

Harriss, D.J., Atkinson, G., Batterham, A., George, K., Cable, N.T., Reilly, T., Haboubi, N., and Renehan, A.G. (2009). Lifestyle factors and colorectal cancer risk (2): a systematic review and meta-analysis of associations with leisure-time physical activity. *Colorectal Disease*, **11(7)**, 689-701.

Haskell, W.L. (1994). J.B. Wolffe Memorial Lecture. Health consequences of physical activity: understanding and challenges regarding dose-response. *Medicine and Science in Sports and Exercise*, **26(6)**, 649-660.

Haskell, W.L. (1984). Physical activity and health: the need to define the required stimulus. *American Journal of Cardiology*, **55**, 4D-9D.

Haskell, W.L., Lee, I.M., Pate, R.R., Powell, K.E., Blair, S.N., Franklin, B.A., Macera, C.A., Heath, G.W., Thompson, P.D., and Bauman, A. (2007). Physical activity and public health: updated recommendation for adults from the American College of Sports Medicine and the American Heart Association. *Circulation*, **116**, 1081-1093.

Hatori, M., Hasegawa, A., Adachi, H., Shinozaki, A., Hayashi, R., Okano, H., Mizunuma, H., and Murata, K. (1993). The effects of walking at the anaerobic threshold level on vertebral bone loss in postmenopausal women. *Calcified Tissue International*, **52(6)**, 411-414.

Healy, G.N., Matthews, C.E., Dunstan, D.W., Winkler, E.A.H., and Owen, N. (2011). Sedentary time and cardio-metabolic biomarkers in US adults: NHANES 2003-06. *European Heart Journal*, **32**, 590-597.

Healy, G.N., Dunstan, D.W., Salmon, J., Cerin, E., Shaw, J.E., Zimmet, P.Z., and Owen, N. (2008). Breaks in sedentary time: beneficial associations with metabolic risk. *Diabetes Care*, **31(4)**, 661-666.

Healy, G.N., Wijndaele, K., Dunstan, D.W., Shaw, J.E., Salmon, J. *et al.* (2008). Objectively measured sedentary time, physical activity and metabolic risk. *Diabetes Care*, **31**, 369-371.

Heikes, K.E., Eddy, D.M., Arondekar, B., and Schlessinger, L. (2008). Diabetes Risk Calculator: A simple tool for detecting undiagnosed diabetes and pre-diabetes. *Diabetes Care*, **31(5)**, 1040-1045.

Hellénus, M-L. and Sundberg, C.J. (2011). Physical activity as medicine: time to translate evidence into clinical practice. *British Journal of Sports Medicine*, **45**, 158 doi:10.1136/bsjm.2011.084244.

Helmerhorst, H.J.F., Wijndaele, K., Brage, S., Wareham, N.J., and Ekelund, U. (2009). Objectively measured sedentary time may predict insulin resistance independent of moderate- and vigorous-intensity physical activity. *Diabetes*, **58**, 1776-1779.

Helmrich, S.P., Ragland, D.R., Leung, R.W., and Paffenbarger, R.S.Jr. (1991). Physical activity and reduced occurrence of non-insulin-dependent diabetes mellitus. *New England Journal of Medicine*, **325**, 147-152.

Henson, J., Yates, T., Biddle, S.J.H., Edwardson, C.L., Khunti, K., Wilmot, E.G., Gray, L.J., Gorely, T., Nimmo, M.A., and Davies, M.J. (2013). Associations of objectively measured sedentary behaviour and physical activity with markers of cardiometabolic health. *Diabetologia*, doi:10.1007/s00125-013-2845-9.

Herman, B., Schmitz, P.I., Leyten, A.C., Van Luijk, J.H., Frenken, C.W., Op De Coul, A.A., and Schulte, B.P. (1983). Multivariate logistic analysis of risk factors for stroke in Tilburg, The Netherlands. *American Journal of Epidemiology*, **118(4)**, 514-525.

Heyward, V.H. and Gibson, A.L. (2014). *Advanced fitness assessment and exercise prescription*, 7th edition. Human Kinetics: Champaign, IL.

Holmes, M.D., Chen, W.Y., Feskanich, D., Kroenke, C.H., and Colditz, G.A. (2005). Physical activity and survival after breast cancer diagnosis. *Journal of the American Medical Association*, **293**, 2479-2486.

Hou, L., Ji, B.T., Blair, A., Dai, Q., Gao, Y.T., and Chow, W.H. (2004). Commuting physical activity and risk of colon cancer in Shanghai, China. *American journal of Epidemiology*, **160(9)**, 860-867.

Howley, E.T. (2001). Type of activity: resistance, aerobic and leisure versus occupational physical activity. *Medicine and Science in Sports and Exercise*, **33(6)**, 364-369.

Hu, F.B., Li, T.Y., Colditz, G.A., Willett, W.C., and Manson, J.E. (2003). Television watching and other sedentary behaviours in relation to risk of obesity and type 2 diabetes mellitus in women. *Journal of the American Medical Association*, **289**, 1785-1791.

Hu, F.B., Qiao, Q., Silventoinen, K., Eriksson, J.G., Jousilahti, P., Lindstrom, J., Valle, T.T., Nissinen, A., Tuomilehto, J. (2003). Occupational, commuting and leisure-time physical activity in relation to risk for type 2 diabetes in middle-aged Finnish men and women. *Diabetologica*, **46**, 322-329.

Hu, F.B., Leitzmann, M.F., Stampfer, M.J., Colditz, G.A., Willett, W.C., and Rimm, E.B. (2001). Physical activity and television watching in relation to risk for type 2 diabetes mellitus in men. *Archives of Internal Medicine*, **161(12)**, 1542-1548.

Hu, F.B., Stampfer, M.J., Colditz, G.A., Ascherio, A., Rexrode, K.M., Willett, W.C., and Manson, J.E. (2000). Physical activity and risk of stroke in women. *Journal of the American Medical Association*, **283**(22), 2961-2967.

Hu, F.B., Sigal, R.J., Rich-Edwards, J.W., Colditz, G.A., Solomon, C.G., Willett, W.C., Speizer, F.E., and Manson, J.E. (1999). Walking compared with vigorous physical activity and risk of type 2 diabetes in women: a prospective study. *Journal of the American Medical Association*, **282**, 1433-1439.

Hunter, G.R., Weinsier, R.L., Bamman, M.M., and Larson, D.E. (1998). A role for high intensity exercise on energy balance and weight control. *International Journal of Obesity*, **22**, 489-493.

Hutchinson, T.M., Whalen, R.T., Cleet, T.M., Vogel, J.M., and Arnaud, S.B. (1995). Factors in daily physical activity related to calcaneal mineral density in men. *Medicine and Science in Sports and Exercise*, **27**. 745-750.

Inoue, M., Iso, H., Yamamoto, S., Kurahashi, N., Iwasaki, M., Sasazuki, S., and Tsugane, S. (2008). Daily total physical activity level and premature death in men and women: Results from a large-scale population-based cohort study in Japan (JPHC Study). *Annals of Epidemiology*, **18**(7), 522-530.

Institute of Medicine (2005). Dietary reference intakes for energy, carbohydrate, fiber, fat, fatty acids, cholesterol, protein, and amino acids. Washington, DC: Institute of Medicine of the National Academics.

International Agency Research Against Cancer (2002). *Weight control, physical activity and cancer*. Chemoprevention: IARC Press.

Irwin, M.L., Yasui, Y., Ulrich, C.M., Bowen, D., Rudolph, R.E., Schwartz, R.S., Yukawa, M., Aielle, E., Potter, J.D., and McTiernan, A. (2003). Effect of exercise on total and intra-abdominal body fat in postmenopausal women: a randomised controlled trial. *Journal of the American Medical Association*, **289**, 323-330.

Jakicic, J.M., Clark, K., Coleman, E., Foreyt, J., Melanson, E., Volek, J., and Volpe, S.L. (2001). American College of Sports Medicine position stand. Appropriate intervention strategies for weight loss and prevention of weight regain for adults. *Medicine and Science in Sports and Exercise*, **33**(12):2145–2156.

Jakicic, J.M., Wing, R.R., Butler, B.A., and Robertson, R.J. (1995). Prescribing exercise in multiple short bouts versus one continuous bout: effect on adherence, cardiorespiratory fitness and weight loss in overweight women. *International Journal of Obesity*, **19**, 893-901.

Jeffery, R.W., Wing, R.R., Thorson, C., and Burton, L.R. (1998). Use of personal trainers and financial incentives to increase exercise in a behavioural weight-loss program. *Journal of Consulting and Clinical Psychology*, **66**, 777-783.

Jeon, C.Y., Lokken, R.P., Hu, F.B., and van Dam, R.M. (2007). Physical activity of moderate intensity and risk of type 2 diabetes: a systematic review. *Diabetes Care*, **30**(3), 550-555.

- Johannsen, D.L., Welk, G.J., Sharp, R.L., and Flakoll, P.J. (2008). Differences in daily energy expenditure in lean and obese women: the role of posture allocation. *Obesity (Silver Spring)*, **16**, 34-39.
- Kahn, S.R., Shrier, I. And Kearon, C. (2008). Physical activity in patients with deep venous thrombosis: A systematic review. *Thrombosis Research*, **122(6)**, 763-773.
- Kallings, L.V., Sierra Johnson, J., Fisher, R.M., Faire, Ud., Ståhle, A., Hemmingsson, E., and Hellénus, M.L. (2009). Beneficial effects of individualised physical activity on prescription on body composition and cardiometabolic risk factors: results from a randomised controlled trial. *European Journal of Cardiovascular Prevention and Rehabilitation*, **16(1)**, 80-84.
- Kang, J., Robertson, R.J., Hagberg, J.M., Kelley, D.E., Goss, F.L., DaSilva, S.G. Suminski, R.R., and Utter, A.C. (1996). Effect of exercise intensity on glucose and insulin metabolism in obese individuals and obese NIDDM patients. *Diabetes Care*, **19(4)**, 341-349.
- Karlsson, M.K., Hasserijs, R., and Obrant, K.J. (1996). Bone mineral density and muscle strength in young male adults with different exercise level. *Calcified Tissue International*, **64**, 490-498.
- Karlsson, M.K., Johnell, O., and Obrant, K.J. (1993). Bone mineral density in weight lifters. *Calcified Tissue International*, **52**, 212-215.
- Katzmarzyk, P.T. (2010). Physical activity, sedentary behaviour, and Health: Paradigm paralysis or paradigm shift? *Diabetes*, **59**, 2717-2725.
- Katzmarzyk, P.T., Church, T.S., Craig, C.L., and Bouchard, C. (2009). Sitting time and mortality from all causes, cardiovascular disease, and cancer. *Medicine and Science in Sports and Exercise*, **41**, 998-1005.
- Katzmarzyk, P.T. and Lee, I-M. (2012). Sedentary behaviour and life expectancy in the USA: a cause-deleted life table analysis. *BMJ Open*, **2**:e000828. doi:10.1136/bmjopen-2012-000828.
- Kelley, G.A. (1999). Aerobic exercise and resting blood pressure among women: a meta-analysis. *Preventive medicine*, **28(3)**, 264-275.
- Kelley, G.A. and Kelley, K.S. (2006). Aerobic exercise and lipids and lipoproteins in men: a meta-analysis of randomised controlled trials. *Journal of Men's Health and Gender*, **31(1)**, 61-70.
- Kelley, G.A. and Sharpe, K.K. (2001). Aerobic exercise and resting blood pressure in older adults: a meta-analytic review of randomised controlled trials. *Journal of Gerontology: Series A, Biological Sciences and Medical Sciences*, **56(5)**, M298-M303.
- Kelley, G.A., Kelley, K.A. and Tran, Z.V. (2001). Aerobic exercise and resting blood pressure: a meta-analysis of randomised controlled trials. *Preventive Cardiology*, **4(2)**, 73-80.
- Kelley, G.A., Kelley, K.A. and Tran, Z.V. (2000). Exercise and bone mineral density in men: a meta-analysis. *Journal of Applied Physiology*, **88**1730-1736.

Kesäniemi, A., Riddoch, C.J., Reeder, B., Blair, S.N., and Sørensen, T.I.A. (2010) Advancing the future of physical activity guidelines in Canada: an independent expert panel interpretation of the evidence. *International Journal of Behavioural Nutrition and Physical Activity*, **7**, 41.

Keum, N., Ju, W., Lee, D.H., Ding, E.L., Hsieh, C.C., Goodman, J.E., and Giovannucci, E.L. (2014). Leisure-time physical activity and endometrial cancer risk: Dose-response meta-analysis of epidemiological studies. *International Journal of Cancer*, **135**, 682-694.

Kiely, D.K., Wolf, P.A., Cupples, L.A., Belser, A.S., and Kannel, W.B., (1994). Physical activity and stroke risk: The Framingham study. *American Journal of Epidemiology*, **140**, 608-620.

Kikkinos P.F., Holland, J.C., Narayan, P., Collieran, J.A., Dotson, C.O., and Papademetriou, V. (1995a). Miles run per week and high density lipoprotein cholesterol levels in healthy, middle-aged men. A dose-response relationship. *Archives of Internal Medicine*, **155(4)**, 415-20.

Kikkinos P.F., Holland, J.C., Pittaras, A.E., Narayan, P., Dotson, C.O., and Papademetriou, V. (1995b). Cardiorespiratory fitness and coronary heart disease risk factor association in women. *Journal of the American College of Cardiology*, **26(2)**, 358-64.

Kikkinos P.F. and Fernhall, B. (1999). Physical activity and high density lipoprotein cholesterol levels. *Sports Medicine*, **28(5)**, 307-14.

King, A.C., Haskell, W.L., Young, D.R., Oka, R.K., and Stefanik, M.L. (1995). Long-term effects of varying intensities and formats of physical activity on participation rates, fitness, and lipoproteins in men and women aged 50 to 65 years. *Circulation*, **91**, 2596-2604.

King, A.C., Haskell, W.L., Taylor, C.B., Kraemer, H.C., and Debusk, R. (1991). Group vs. Home-based exercise training in healthy older men and women: a community-based clinical trial. *Journal of the American Medical Association*, **266**, 1535-1542.

Kingwell, B.A. and Jennings, G. (1993). Effects of walking and other exercise programs upon blood pressure in normal subjects. *Medical journal of Australia*, **158**, 234-238.

Knols, R., Aaronson, N.K., Uebelhart, D., Fransen, J., and Aufdemkampe, G. (2005) Physical exercise in cancer patients during and after medical treatment: a systematic review of randomised and controlled clinical trials. *Journal of Clinical Oncology*, **23**, 3830-3842.

Knowler, W.C., Barrett-Connor, E., Fowler, S.E., Hamman, R.F., Lachin, J.M., Walker, E.A., and Nathan, D.M. (2002). Reduction in the incidence of type 2 diabetes with lifestyle intervention or metformin. *New England Journal of Medicine*, **346**, 393-403.

Kodama, S., Tanaka, S., Saito, K., Shu, M., Sone, Y., Onitake, F., Suzuki, E., Shimano, H., Yamamoto, S., Kondo, K., Ohashi, Y., Yamada, N., and Sone, H. (2007). Effect of aerobic exercise training on serum levels of high-density lipoprotein cholesterol: a meta-analysis. *Archives of Internal Medicine*, **167(10)**, 999-1008.

Kokkinos, P. and Myers, J. (2010). Exercise and physical activity: Clinical outcomes and applications. *Circulation*, **122**, 1637-1648.

Koster, A., Caserotti, P., Patel, K.V., Matthews, C.E., Berrigan, D., Van Domelen, D.R., Brychta, R.J., Chen, K.Y., and Harris, T.B. (2012). Association of sedentary time with mortality independent of moderate to vigorous physical activity. *PLoS ONE*, **7(6)**: e37696. doi:10.1371/journal.pone.0037696.

Kramer, A.F., Hahn, S., Cohen, N.J., Banich, M.T., McAuley, E., Harrison, C.R., Chason, J., Vakil, E., Bardell, L., Boileau, R.A., and Colcombe, A. (1999). Ageing, fitness and neurocognitive function. *Nature*, **400**, 418-419.

Kraus, W.E., Houmard, J.A., Duscha, B.D., Knetzger, K.J., Wharton, M.B., McCartney, J.S., Bales, C.W., Henes, S., Samsa, G.P., Otvos, J.D., Kulkarni, K.R., and Slentz, C.A. (2002). Effects of the amount and intensity of exercise on plasma lipoproteins. *New England Journal of Medicine*, **347**, 1483-1492.

Kujala, U.M. (2009). Evidence on the effects of exercise therapy in the treatment of chronic disease. *British Journal of Sports Medicine*, **43**, 550-555.

Laaksonen, D.E., Linström, J., Lakka, T.A., Eriksson, J.G., Niskanen, L., Wikström, K., Aunola, S., Keinänen-Kiukaanniemi, S., Laakso, M., Valle, T.T., Ilanne-Parikka, P., Louheranta, A., Hämäläinen, H., Rastas, M., Salminen, V., Cepaitis, Z., Hakumäki, M., Kaikkonen, H., Härkönen, P., Sundvall, J., Tuomilehto, J., and Uusitupa, M. (2005). Physical activity in the prevention of type 2 diabetes: the Finnish diabetes prevention study. *Diabetes*, **54(1)**, 158-165.

Lagerros, Y.T., Hsieh, S.F. and Hsieh, C.C. (2004). Physical activity in adolescence and young adulthood and breast cancer risk: a quantitative review. *European Journal of Cancer Prevention*, **13(1)**, 5-12.

Lahmann, P.H., Friedenreich, C., Schuit, A.J., Salvini, S., Allen, N.E., Key, T.J., Khaw, K.T., Bingham, S., Peeters, P.H., Monninkhof, E., Bueno-de-Mesquita, H.B., Wirfält, E., Manjer, J., Gonzales, C.A., Ardanaz, E., Amiano, P., Quirós, J.R., Navarro, C., Martinez, C., Berrino, F., Palli, D., Tumino, R., Panico, S., Vineis, P., Trichopoulou, A., Bamia, C., Trichopoulos, D., Boeing, H., Schulz, M., Linseisen, J., Chang-Claude, J., Chapelon, F.C., Fournier, A., Boutron-Ruault, M.C., Tjønneland, A., Føns Johnson, N., Overvad, K., Kaaks, R., Riboli, E. (2007). Physical activity and breast cancer risk: the European prospective investigation into cancer and nutrition. *Cancer Epidemiology, Biomarkers and Prevention*, **16(1)**, 36-42.

Larsen, R.N., Kingwell, B.A., Sethi, P., Cerin, E., Owen, N., and Dunstan, D.W. (2014) Breaking up prolonged sitting reduces resting blood pressure in overweight/obese adults. *Nutrition, Metabolism and Cardiovascular Diseases: NMCD*, **24(9)**, 976-982.

Larson, E.B., Wang, L., Bowen, J.D., McCormick, W.C., Teri, L., Crane, P., and Kukull, W. (2006). Exercise is associated with reduced risk for incident dementia among persons 65 years of age and older. *Annals of Internal Medicine*, **144(2)**, 73-81.

Latouche, C., Jowett, J.B., Carey, A.L., Bertovic, D.A., Owen, N., Dunstan, D.W., and Kingwell, B.A. (2013). Effects of breaking up prolonged sitting on skeletal muscle gene expression. *Journal of Applied Physiology*, **114(4)**, 453-460.

Laukkanen, J.A., Rauramaa, R., Mäkitallio, T.H., Toriola, A.T., and Kurl, S. (2011). Intensity of leisure-time physical activity and cancer mortality in men. *British Journal of Sports Medicine*, **45(2)**, 125-129.

Lautenschlager, N.T., Cox, K., and Cyarto, E.V. (2012). The influence of exercise on brain aging and dementia. *Biochimica et Biophysica Acta*, **1822**, 474-481.

Lavie, C.J., Church, T.S., Milani, R.V., and Ernest, C.P. (2011). Impact of physical activity, cardiorespiratory fitness, and exercise training on markers of inflammation. *Journal of Cardiopulmonary Rehabilitation and Prevention*, **31(3)**, 137-145.

Lawler, P.R., Filion, K.B., and Eisenberg, M.J. (2011). Efficacy of exercise-based cardiac rehabilitation post-myocardial infarction: A systematic review and meta-analysis of randomised controlled trials. *American Heart Journal*, **162(4)**, 571-584.

Lee, C.D., Folsom, A.R. and Blair, S.N. (2003). Physical activity and stroke risk. *Stroke*, **34**, 2475-2481.

Lee, D.C., Sui, X., Ortega, F.B., Kim, Y-S., Church, T.S., Winett, R.A., Ekelund, U., Katzmarzyk, P.T., and Blair, S.N. (2011). Comparisons of leisure-time physical activity and cardiorespiratory fitness as predictors of all cause mortality in men and women. *British Journal of Sports Medicine*, **45**, 504-510.

Lee, I.M. (2008). "Dose-Response Relation between Physical Activity and Health: An Observational Epidemiologic Prospective." Available from: www.pitt.edu/~super7/31011-32001/31521.ppt

Lee, I-M., Shiroma, E.J., Lobelo, F., Puska, P., Blair, S.N., and Katamarzyk, P.T. (2012). Effect of physical inactivity on major non-communicable disease worldwide: an analysis of burden of disease and life expectancy. *The Lancet*, **380**, 219-229.

Lee, I.M., Djoussé, L., Sesso, H.D., Wang, L., and Buring, J.E. (2010). Physical activity and weight gain prevention. *Journal of the American Medical Association*, **303(12)**, 1173-1179.

Lee, I.M., Sesso, H.D., Oguma, Y., and Paffenbarger, R.S. Jr. (2004). The "weekend warrior" and risk of mortality. *American Journal of Epidemiology*, **160(7)**, 636-641.

Lee, I.M., Rexrode, K.M., Cook, N.R., Heenekens, C.H., and Burin, J.E. (2001). Physical activity and breast cancer risk: the Women's health Study (United States). *Cancer Causes and Control*, **12(2)**, 137-145.

Lee, I.M. and Skerrett, P.J., (2001). Physical activity and all-cause mortality: what is the dose-response relation? *Medicine and Science in Sports and Exercise*, **33(6 Suppl.)**, S459-S471; discussion S493-S494.

Lee, I.M. and Paffenbarger, R.S. Jr. (1998) Physical Activity and Stroke Incidence. *Stroke*, **29**, 2049-2054.

Lee, I-M., Hsieh, C-C., and Paffenbarger, R.S. (1995). Exercise intensity and longevity in men. The Harvard Alumni Study. *Journal of the American Medical Association*, **73**, 1179-1184.

Lee, I-M., Paffenbarger, R.S. and Hsieh, C-C. (1992). Time trends in physical activity among college alumni, 1962-1988. *American Journal of Epidemiology*, **135(8)**, 915-925.

Lee, I.M., Paffenbarger, R.S. Jr. and Hsieh, C. (1991). Physical activity and risk of developing colorectal cancer among college alumni. *Journal of the National Cancer Institute*, **83(18)**, 1324-1329.

Lee, K.J, Inoue, M., Otani, T., Iwasaki, M., Sasazuki, S, and Tsugane, S. (2007). Physical activity and risk of colorectal cancer in Japanese men and women: the Japan Public Health Center-based prospective study. *Cancer Causes and Control*, **18(2)**, 199-209.

Lee, S.M.C., Moore, A.D., Everett, M.E., Stenger, M.B., Wareham, N.J., and Ekelund, U. (2009). Aerobic exercise deconditioning and countermeasures during bed rest. *Aviation, Space, and Environmental Medicine*, **81**, 52-63.

Leichter, I., Simkin, A., Margulies, J.Y., Bivas, A., Steinberg, R., Giladi, M., and Milgrom, C. (1989). Gain in mass density of bone following strenuous physical activity. *Journal of Orthopaedic Research*, **7**, 86-90.

Levi, F., La Vecchia, C., Negri, E., and Franceschi, S. (1993). Selected physical activities and the risk of endometrial cancer. *British Journal of Cancer*, **67**, 846-851.

Levine, J.A., Lanningham-Foster, L.M., McCrady, S.K., et al. (2005). Interindividual variation in posture allocation: possible role in human obesity. *Science*, **307**, 584-586.

Levine, J.A. (2003). Non exercise activity thermogenesis. *Proceedings of the Nutrition Society*, **62(3)**, 667-679.

Li, J. and Siegrist, J. (2012). Physical activity and risk of cardiovascular disease – a meta-analysis of prospective cohort studies. *International Journal of Environmental Research and Public Health*, **9(2)**, 391-407.

Linström, J. and Tuomilehto, J. (2003). The Diabetes Risk Score: A practical tool to predict type 2 diabetes risk. *Diabetes Care*, **26(3)**, 725-731.

Linström, J., Ilanne-Parikka, P., Peltonen, M., Aunola, S., Eriksson, J.G., Hemiö, K., Hämäläinen, H., Härkönen, P., Keinänen-Kiukaanniemi, S., Laakso, M., Louheranta, A., Mannelin, M., Paturi, M., Sundvall, J., Valle, T.T., Uusitupa, M., and Tuomilehto, J. (2006). Sustained reduction in the incidence of type 2 diabetes by lifestyle intervention: follow-up of the Finnish Diabetes Prevention Study. *Lancet*, **368(9548)**, 1673-1679.

Lipman, R.L., Raskin, P., Love, T., Triebwasser, J., Lecocq, F.R., and Schnure, J.J. (1972). Glucose intolerance during decreased physical activity in man. *Diabetes*, **21**, 101-107.

Löllgen, H., Böckenhoff, A. and Knapp, G. (2009) Physical activity and all-cause mortality: an updated meta-analysis with different intensity categories. *International Journal of Sports Medicine*, **30(3)**, 213-224.

Louis, E.D., Tang, M.X., Schupf, N., and Mayeux, R. (2005). Functional correlates and prevalence of mild parkinsonian signs in a community population of older people. *Archives of Neurology*, **62**, 297-302.

Lyden, K., Kozey Keadle, S.L., Staudenmayer, J.W., and Freedson, P.S. (2012). Validity of two wearable monitors to estimate breaks from sedentary time. *Medicine and Science in Sports and Exercise*, **44(11)**, 2243-2252.

Lynch, J., Helmrish, S.P., Lakka, T.A., Kaplan, G.A., Cohen, R.D., Salonen, R., and Salonen, J.T. (1996). Moderately intense physical activities and high levels of cardiorespiratory fitness reduce the risk of non-insulin-dependent diabetes mellitus in middle-aged men. *Archives of Internal Medicine*, **156**, 1307-1314.

MacDougall, J.D., Webber, C.E., Martin, J., Ormerod, S., Chesley, A., Younglai, E.V., Gordon, C.L., and Blimkie, C.J. (1992). Relationship among running mileage, bone density, and serum testosterone in male runners. *Journal of Applied Physiology*, **73(3)**, 1165-1170.

Madsen, K.L., Adams, W.C., and Van Loan, M.D. (1998). Effects of physical activity, body weight and composition, and muscular strength on bone density in young women. *Medicine and Science in Sports and Exercise*, **30**, 114-120.

Mai, P.L., Sullivan-Halley, J., Ursin, G., Stram, D.O., Deapen, D., Villaluna, D., Horn-Ross, P.L., Clarke, C.A., Reynolds, P., Ross, R.K., West, D.W., Anton-Culver, H., Ziogas, A., and Bernstein, L. (2007). Physical activity and colon cancer risk among women in the California Teachers Study. *Cancer Epidemiology, Biomarkers and Prevention*, **16(3)**, 517-525.

Manson, J.E., Greenland, P., LaCroix, A.Z., Stefanick, M.L., Mouton, C.P., Oberman, A., Perri, M.G., Sheps, D.S., Pettinger, M.B., and Siscovick, D.S. (2002). Walking compared with vigorous exercise for the prevention of cardiovascular events in women. *New England Journal of Medicine*, **347(10)**, 716-725.

Manson, J.E., Hu, F.B., Rich-Edwards, J.W., Colditz, G.A., Stampfer, M.J., Willett, W.C., Speizer, F.E., and Hennekens, C.H. (1999). A prospective study of walking compared with vigorous exercise in the prevention of coronary heart disease in women. *New England Journal of Medicine*, **341**, 650-658.

Manson, J.E., Nathan, D.M., Krolewski, A.S., Stampfer, M.J., Willett, W.C., and Hennekens, C.H. (1992). A prospective study of exercise and incidence of diabetes among US male physicians. *Journal of the American Medical Association*, **268**, 63-67.

Manson, J.E., Stampfer, M.J., Colditz, G.A., Willett, W.C., Rosner, B., Hennekens, C.H., Speizer, F.E., Rimm, E.B., and Krolewski, A.S. (1991). Physical activity and incidence of non-insulin dependent diabetes mellitus in women. *Lancet*, **338**, 774-778.

Marcus, P.M., Newcomb, P.A., Moorman, P.G., Millikan, R.C., Baird, D.D., Qagish, B., and Sternfeld, B. (1999). Physical activity at age 12 and adult breast cancer risk (United States). *Cancer Causes and Control*, **10(4)**, 293-302.

Martinez, M.E., Giovannucci, E., Spiegelman, D., Hunter, D.J., Willett, W.C., and Colditz, G.A. (1997). Leisure-time physical activity, body size, and colon cancer in women. Nurses' Health Study Research Group. *Journal of the National Cancer Institute*, **89(13)**, 948-955.

Matthews, C.E., George, S.M., Moore, S.C., Bowles, H.R., Blair, A., Park, Y., Troiano, R.P., Hollenbeck, A., and Schatzkin, A. (2012). Amount of time spent in sedentary behaviours and cause-specific mortality in US adults. *American Journal of Clinical Nutrition*, **95(2)**, 437-445.

- Matthews, C.E., Chen, K.Y., Freedson, P.S., Buchowski, M.S., Beech, B.M., Pate, R.R., and Troiano, R.P. (2008). Amount of time spent in sedentary behaviours in the United States, 2003-2004. *American Journal of Epidemiology*, **167**(7), 875-881.
- Matusaki, M., Ikeda, M., Tashiro, E., Koga, M., Miura, S., Ideishi, M., Tanaka, H., Shindo, M., and Arakawa, K. (1992). Influence of workload on the antihypertensive effect of exercise. *Clinical and Experimental Pharmacology and Physiology*, **19**(7), 471-479.
- Mayoux-Benhamou, M.A., Leyge, J.F., Roux, C., and Revel, M. (1999). Cross-sectional study of weight-bearing activity on proximal femur bone mineral density. *Calcified Tissue International*, **64**, 179-183.
- McCrady, S.K. and Levine, J.A. (2009). Sedentariness at work; how much do we really sit? *Obesity (Silver Spring)*, **17**(11), 2103-2105.
- McTiernan, A., Sorensen, B., Irwin, M.L., Morgan, A., Yasui, Y., Rudolph, R.E., Surawicz, C., Lampe, J.W., Lampe, P.D., Ayub, K., and Potter, J.D. (2007). Exercise effect on weight and body fat in men and women. *Obesity*, **15**, 1496-1512.
- Mekary, R.A., Feskanich, D., Malspeis, S., Hu, F.B., Willet, W.C., and Field, A.E. (2009). Physical activity patterns and prevention of weight gain in premenopausal women. *International Journal of Obesity*, **33**(9), 1039-1047.
- Metcalfe, R., Babraj, J., Fawcner, S., and Volvaard, N. (2012). Towards the minimal amount of exercise for improving metabolic health: Beneficial effects of reduced-exertion high-intensity interval training. *European Journal of Applied Physiology*, **112**(7), 2767-2775.
- Miller, J.P., Pratley, R.E., Goldberg, A.P., Rubin, M., Treuth, M.S., Ryan, A.S., and Hurley, B.F. (1994). Strength training increases insulin action in healthy 50- to 65-yr-old men. *Journal of Applied Physiology*, **77**, 1122-1127.
- Mittelman, M.A., Maclure, M., Tofler, G.H., Sherwood, J.B., Goldberg, R.J., and Muller, J.E. (1993). Triggering of acute myocardial infarction by heavy physical exertion: protection against triggering by regular exertion. *New England Journal of Medicine*, **329**, 1677-1683.
- Molz, A.B., Heyduck, B., Lill, H., Spanutii, E., and Rocker, L. (1993). The effect of different exercise intensities on the fibrolytic system. *European Journal of Applied Physiology*, **67**, 298-304.
- Monninkhof, E.M., Elias, S.G., Vlems, F.A., van der Twel, I., Schuit, A.J., Voskuil, D.W., and van Leeuwen, F.E. (2007). Physical activity and breast cancer: a systematic review. *Epidemiology*, **18**(1), 137-157.
- Moreira, W.D., Fuchs, F.D., Ribeiro, J.P., and Appel, L.J. (1999). The effects of two aerobic training intensities on ambulatory blood pressure in hypertensive patients: results of a randomised trial. *Journal of Clinical Epidemiology*, **52**, 637-642.
- Morris, J.N., and Crawford, M.D. (1958). Coronary heart disease and physical activity of work: evidence of a national necropsy survey. *British Medical Journal*, **2**, 1485-1496.

Morris, J.N. (1994). Exercise in the prevention of coronary heart disease: today's best buy in public health. *Medicine and Science in Sports and Exercise*, **26**, 807-814.

Murray, C.J.L., Richards, M.A., Newton, J.N., Fenton, K.A., Anderson, H.R., Atkinson, C., Bennett, D., Bernabé, E., Blencowe, H., Bourne, R., Braithwaite, T., Brayne, C., Bruce, N.G., Brugha, T.S., Burney, P., Dherani, M., Dolk, H., Edmond, K., Ezzati, M., Flaxman, A.D., Fleming, T.D., Freedman, G., Gunnell, D., Hay, R.J., Hutchings, S.J., Lockett Ohno, S., Lozano, R., Lyons, R.A., Marcenés, W., Naghavi, M., Newton, R., Pearce, N., Pope, D., Rushton, L., Salomon, J.A., Shibuya, K., Vos, T., Wang, H., Williams, H.C., Woolf, A.D., Lopez, A.D., and Davis, A. (2013). UK health performance: findings of the Global Burden of Disease Study 2010. *The Lancet*, **381**, 997-1020.

Murphy, M.H., Blair, S.N. and Murtagh, E.M. (2009). Accumulated versus continuous exercise for health benefit: a review of empirical studies. *Sports Medicine*, **39(1)**, 29-43.

Murphy, M., Nevill, A., Neville, C., Biddle, S., and Hardman, A. (2002). Accumulating brisk walking for fitness, cardiovascular risk, and psychological health. *Medicine and Science in Sports and Exercise*, **34**, 1468-1474.

Murphy, M.H., Nevill, A.M., Murtagh, E.M., and Holder, R.L. (2007). The effect of walking on fitness, fatness and resting blood pressure: a meta-analysis of randomised, controlled trials. *Preventive Medicine*, **44(5)**, 377-385.

Myers, J., Prakesh, M., Froelicher, V., Do, D., Partington, S., and Atwood, J.E. (2002). Exercise capacity and mortality among men referred for exercise testing. *New England Journal of Medicine*, **346**, 793-801.

Need, A.G., Wishard, J.M., Scopacasa, F., Horowitz, M., Morris, H.A., and Nordin, B.E. (1995). Effect of physical activity on femoral bone density on men. *British Medical Journal*, **310**, 1501-1502.

Neilson, H.K., Friedenreich, C.M., Brockton, N.T., and Millikan, R.C. (2009). Physical activity and postmenopausal breast cancer: proposed biologic mechanisms and areas for future research. *Cancer Epidemiology, Biomarkers and Prevention*, **18(1)**, 11-27.

Nikander, R., Sievänen, H., Heinonen, A., Daly, R.M., Uusi-Rasi, K., and Kannus, P. (2010). Targeted exercise against osteoporosis: A systematic review and meta-analysis for optimising bone strength throughout life. *BMC Medicine*, **8**, 47 doi:10.1186/1741-7015-8-47.

Nikander, R., Kannus, P., Dastidar, P., Hannula, M., Harrison, L., Cervinka, T., Narra, N.G., Aktour, R., Arola, T., Eskola, H., Soimakallio, S., Heinonen, A., Hyttinen, J., and Sievänen, H. (2009). Targeted exercises against hip fragility. *Osteoporosis International*, **20**, 1321-1328.

Nocon, M., Hiemann, T., Muller-Riemenschneider, F., Thalau, F., Roll, S., and Willich, S.N. (2008). Association of physical activity with all-cause and cardiovascular mortality: A systematic review and meta-analysis. *European Journal of Cardiovascular Prevention and Rehabilitation*, **15**, 239-246.

O'Donovan, G., Blazeovich, A.J., Boreham, C., Cooper, A.R., Crank, H., Ekelund, U., Fox, K.R., Gately, P., Giles-Corti, B., Gill, J.M.R., Hamer, M., McDermott, I., Murphy, M., Mutrie, N., Reilly, J.J., Saxton, J.M., and Stamatakis, E. (2010). The ABC of Physical Activity for health: A consensus statement from the British Association of Sport and Exercise Sciences. *Journal of Sport Sciences*, **28(6)**, 573-591.

Okada, K., Hayashi, T., Tsumura, K., Suematsu, C., Endo, G., and Fujii, S. (2000). Leisure-time physical activity at weekends and the risk of type 2 diabetes mellitus in Japanese men: the Osaka Health Survey. *Diabetic Medicine*, **17**, 53-58.

Olsen, C.M., Green, A.C., Whiteman, D.C., Sedeghi, S., Kolahdooz, F. And Webb, P.M. (2007). Obesity and the risk of epithelial ovarian cancer: a systematic review and meta-analysis. *European Journal of Cancer*, **43(4)**, 690-709.

Owen, N., Healy, G.N., Matthews, C.E., and Dunstan, D.W. (2010). Too much sitting: the population health science of sedentary behaviour. *Exercise and Sport Sciences Reviews*, **38**, 105-113.

Paffenbarger, R.S. Jr., Lee, I-M. and Leung, R. (1994) Physical activity and personal characteristics associated with depression and suicide in American college men. *Acta Psychiatrica Scandinavica Supplementum*, **377**, 16-22.

Paffenbarger, R.S.Jr., Hyde R.T., Wing, A.L., Lee, I.M., Jund, D.L., and Kampert, J.B. (1993). The association of changes in physical-activity level and other lifestyle characteristics with mortality among men. *New England Journal of Medicine*, **328**, 538-545.

Paffenbarger, R.S.Jr., Hyde, R.T., Wing, A.L., and Hsieh, C.C. (1986). Physical activity, all-cause mortality, and longevity of college alumni. *New England Journal of Medicine*, **314**, 605-613.

Paffenbarger, R.S.Jr., Hyde R.T., Wing, A.L., and Steinmetz, C.H. (1984). A natural history of athleticism and cardiovascular health. *Journal of the American Medical Association*, **252**, 491-495.

Paffenbarger, R.S.Jr., Wing, A.L., Hyde R.T., and Jung, D.L. (1983). Physical activity and the incidence of hypertension in college alumni. *American Journal of Epidemiology*, **117**, 245-257.

Paffenbarger, R.S.Jr., Wing, A.L. and Hyde R.T. (1978). Physical activity as an index of heart attack risk in college alumni. *American Journal of Epidemiology*, **108**, 161-175.

Pahor, M., Guralnik, J.M., Ambrosius, W.T., Blair, S., Bonds, D.E., Church, T.S., Espeland, M.A., Fielding, R.A., Gill, T.M., Groessl, E.J., King, A.C., Kritchevsky, S.B., Manini, T.M., McDermott, M.M., Miller, M.E., Newman, A.B., Rejeski, W.J., Sink, K.M., and Williamson, J.D. (2014). Effect of structured physical activity on prevention of major mobility disability in older adults. *Journal of the American Medical Association*, **311(23)**, 2387-2396.

Pan, X.R., Li, G.W., Hu, Y.H., Wang, J.X., Yang, W.Y., An, Z.X., Hu, Z.X., Lin, J., Xiao, J.Z., Cao, H.B., Liu, P.A., Jiang, X.G., Jiang, Y.Y., Wang, J.P., Zheng, H., Zhang, H., Bennett, P.H., and howard, B.V. (1997). Effects of diet and exercise in preventing NIDDM in people with impaired glucose tolerance. The Da Qing IGT and Diabetes Study. *Diabetes Care*, **20(4)**, 537-544.

Pandey, A., Garg, S., Khunger, M., Ayers, C., Kumbhani, D., Levine, B., Deleemos, J., and Berry, J. (2015). Dose response relationship between physical activity and risk of heart failure: A meta-analysis. *Circulation*, **131**, AMP02.

Parry, S. and Straker, L. (2013). The contribution of office work to sedentary behaviour associated risk. *BMC Public Health*, **13**, 296.

Peddie, M.C., Bone, J.L., Rehrer, N.J., Skeaff, C.M., Gray, A.R., and Perry, T.L. (2013). Breaking prolonged sitting reduces postprandial glycemia in healthy normal-weight adults: a randomised crossover trial. *American Journal of Clinical Nutrition*, doi: **10.3495/ajcn.112.051763**

Pescatello, L.S., Guidry, M.A., Blanchard, B.E., Kerr, A., Taylor, A.L., Johnson, A.N., Maresh, C.M., Rodriguez, N., and Thompson, P.D. (2004). Exercise intensity alters postexercise hypertension. *Journal of Hypertension*, **22**, 1881-1888.

Peterson, R.C. (2004). Mild cognitive impairment as a diagnostic entity. *Journal of Internal Medicine*, **256(3)**, 183-194.

Pettersson, U., Nordstrom, P. and Lorentzon, R. (1999). A comparison of bone mineral density in athletes during and after career: A comparison between loaded and unloaded skeletal regions. *Calcified Tissue International*, **59**, 245-248.

Pharm, N.M., Mizoue, T., Tanaka, K., Tsuji, I., Tamakoshi, A., Matsuo, K., Ito, H., Wakai, K., Nagata, C., Sasazuki, S., Inoue, M., and Tsugane, S. (2012). Physical activity and colorectal cancer risk: an evaluation based on a systematic review of epidemiologic evidence among the Japanese population. *Japanese Journal of Clinical Oncology*, **42(1)**, 2-13.

Pikula, A., Beiser, A.S., Chen, T.C., Preis, S.R., Vargias, D., DeCarli, C., Au, R., Kelly-Hayes, M., Kase, C.S., Wolf, P.A., Vasan, R.S., and Seshadri, S. (2013). Serum brain-derived neurotrophic factor and vascular endothelial growth factor levels are associated with risk of stroke and vascular brain injury: Framingham study. *Stroke*, **44(10)**, 2768-2775.

Podewils, L.J., Guallar, E., Kuller, L.H., Fried, L.P., Lopez, O.L., Carlson, M., and Lyketsos, C.G. (2005). Physical activity, APOE genotype, and dementia risk: findings from the Cardiovascular Health Cognition Study. *American Journal of Epidemiology*, **161(7)**, 639-651.

Pollock, M.L., Feigenbaum, M.S. and Brechue, W.F. (1995). Exercise prescription for physical fitness. *Quest*, **47**, 320-337.

Pollock, M.L., Gaesser, G.A., Butcher, J.D., Despres, J.P., Dishman, R.K., Franklin, B.A., and Ewing-Garber, C. (1998). American College of Sports Medicine Position Stand on the recommended quantity and quality of exercise for developing and maintaining cardiorespiratory and muscular fitness, and flexibility in healthy adults. *Medicine and Science in Sports and Exercise*, **30**, 975-991.

Powell, K.E., Thompson, P.D., Caspersen, C.J. and Kendrick, J.S. (1987). Physical activity and the incidence of coronary artery disease. *Annual Review of Public Health*, **8**, 253-287.

President's Council on Fitness, Sports and Nutrition (2012). Too much sitting: Health risks of sedentary behaviour and opportunities for change. *Research Digest*, **13(3)**.

Rakobowchuk, M., Tanguay, S., Burgomaster, K.A., Howarth, K.R., Gibala, M.J., and McDonald, M.J. (2008). Sprint interval and traditional endurance training induce similar improvements in peripheral arterial stiffness and flow-mediated dilation in healthy humans. *American Journal Physiology*, **295**, R236-R242.

Ravussin, E., Lillioja, S., Knowler, W.C., Christin, L., Freymond, D., Abbott, W.G., Boyce, V., Howard, B.V., and Bogardus, C. (1988). Reduced rate of energy expenditure as a risk factor for body-weight gain. *New England Journal of Medicine*, **318**, 467-472.

Ready A.E., Naimark, B., Ducas, J., Sawatzky, J.V., Boreskie, S.L., Drinkwater, D.T., and Oosterveen, S. (1996). Influence of walking volume on health benefits in women post-menopause. *Medicine and Science in Sports and Exercise*, **28(9)**, 1097-1105.

Ribom, E.L. and Piehl-Aulin, K. (2010). Osteoporosis. In: *Physical Activity in the Prevention and Treatment of Disease*. Professionals Associations for Physical Activity, Sweden. Elanders: Sweden.

Rockhill, B., Willett, W.C., Hunter, D.J., Manson, J.E., Hankinson, S.E., Colditz, G.A. (1999). A prospective study of recreational physical activity and breast cancer risk. *Archives of Internal Medicine*, **159(19)**, 2290-2296.

Rogers, M.W., Probst, M.M., Gruber, J.J., Berger, R., and Boone, J.B. (1996). Differential effects of exercise training intensity on blood pressure and cardiovascular responses to stress in borderline hypertensive humans. *Journal of Hypertension*, **14**, 1369-1375.

Roman, O., Camuzzi, A.L., Villalon, E., and Klenner, C. (1981). Physical training program in arterial hypertension. A long-term prospective follow-up. *Cardiology*, **67**, 230-243.

Romijn, J.A., Coyle, E.F., Sidossis, L.S., Gastaldelli, A., Horowitz, J.F., Endert, E., and Wolfe, R.R. (1993). Regulation of endogenous fat and carbohydrate metabolism in relation to exercise intensity and duration. *American Journal of Physiology*, **265(3)**, E380-E391.

Rosengren, A. and Wilhelmsen, L. (1997). Physical activity protects against coronary death and deaths from all causes in middle-aged men: evidence from a 20-year follow-up of the primary prevention study in Goteborg. *Annals of Epidemiology*, **7**, 69-75.

Ross, R., Janssen, I., Dawson, J., Kungl, A., Kuk, J.L., Wong, S.L., Nguyen-Day, T., Lee, A., Kilpatrick, K., and Hudson, R. (2004). Exercise-induced reduction in obesity and insulin resistance in women: a randomised controlled trial. *Obesity Research*, **12**, 789-798.

Ross, R., Dagnone, D., Jones, P.J., Smith, H., Paddags, A., Hudson, R., and Janssen, I. (2000). Reduction in obesity and related comorbid conditions after diet-induced weight loss or exercise-induced weight loss in men. A randomised, controlled trial. *Annals of Internal Medicine*, **133(2)**, 92-103.

Royal College of Physicians (2012). *Exercise for life: physical activity in health and disease*. London: RCP, <http://rcplondon.ac.uk/sites/default/files/documents/exercise-for-life-final.pdf>

Sacco, R.L., Gan, R., Boden-Albala, B., Lin, I.F., Kargman, D.E., Hauser, W.A., Shea, S., and Paik, M.C. (1998). Leisure-time physical activity and ischemic stroke risk: the Northern Manhattan Stroke Study. *Stroke*, **29(2)**, 380-387.

Salmon, J. (2010). Standing up for children's health and education: questioning the sedentary nature of classrooms. The Biannual Fritz Duras lecture, Dean's Lecture Series, University of Melbourne, July 27th, 2010. Samad, A.K.A., Taylor, R.S., Marshall, T., and Chapman, M.A.S. (2005). A meta-analysis of

the association of physical activity with reduced risk of colorectal cancer. *Colorectal Disease*, **7(3)**, 204-2013.

Samad, A.K., Taylor, R.S., Marshall, T., and Chapman, M.A. (2005). A meta-analysis of the association of physical activity with reduced risk of colorectal cancer. *Colorectal Disease*, **7(3)**, 204-213.

Samitz, G., Egger, M., and Zwahlen, M. (2011) Domains of physical activity and all-cause mortality: a systematic review and dose-response meta-analysis of cohort studies. *International Journal of Epidemiology*, **40(5)**, 1382-1400.

Santiago, M.C., Alexander, J.F., Stull, G.A., Serfass, R.C., Hayday, A.M., and Leon, A.S. (1987). Physiological responses of sedentary women to a 20-week conditioning program of walking or jogging. *Scandinavian Journal of Sports Science*, **9**, 33-39.

Saris, W.H., Blair, S.N., van Baak, M.A., Eaton, S.B., Davies, P.S., Di Pietro, L., Fogelholm, M., Rissanen, A., Schoeller, D., Swinburn, B., Tremblay, A., Westerterp, K.R., and Wyatt, H. (2003). How much physical activity is enough to prevent unhealthy weight gain? Outcome of the IASO 1st Stock Conference and consensus statement. *Obesity Reviews*, **4**, 101-114.

Sattelmair, J., Pertman, J., Ding, E.L., Kohl III, H.W., Haskell, W., and Lee, I-M. (2011). Dose response between physical activity and risk of coronary heart disease: A meta-analysis. *Circulation*, **124**, 789-795.

Saunders, T.J., Larouche, R., Colley, R.C., and Tremblay, M.S. (2012). Acute sedentary behaviour and markers of cardiometabolic risk: A systematic review of intervention studies. *Journal of Nutrition and Metabolism*, **Article ID 712435**, 12 pages. Doi:10.1155/2012/712435.

Scheers, T., Philippaerts, R. and Lefevre, J. (2013). Compliance with different physical activity recommendations and its association with socio-demographic characteristics using an objective measure. *BMC Public Health*, **13**, 136 doi:10.1186/1471-2458-13-136.

Schlosser Covell, G.E., Hoffman-Snyder, C.R., Wellik, K.E., Woodruff, B.K., Geda, Y., Caselli, R.J., Demaerschalk, B.M., and Wingerchuk, D.M. (2015). Physical activity level and future risk of mild cognitive impairment or dementia: A critically appraised topic. *Neurologist*, **19(3)**, 89-91.

Schmid, D., Behrens, G., Keimling, M., Jochem, C., Ricci, C., and Leitzmann, M. (2015). A systematic review and meta-analysis of physical activity and endometrial cancer risk. *European Journal of Epidemiology*, **30(5)**, doi: 10.1007/s10654-015-0017-6.

Schoeller, D. (1998). Balancing energy expenditure and body weight. *American journal of Clinical Nutrition*, **68(suppl.)**, 956S-961S.

Schriger, D.L. (2001). Analysing the relationship of exercise and health: methods, assumptions, and limitations. *Medicine and Science in Sports and Exercise*, **33(6)**, 359-363.

Schulze, M.B., Hoffmann, K., Boeing, H., Linseisen, J., Rohrmann, S., Möhlig, M., Pfeiffer, A.F.H., Spranger, J., Thamer, C., Häring, H-U., Fritsche, A., and Joost, H-G. (2007). An accurate risk score based on anthropometric, dietary, and lifestyle factors to predict the development of type 2 diabetes. *Diabetes Care*, **30(3)**, 510-515.

Seguin, R., Buchner, D.M., Liu, J., Allison, M., Manini, T., Wang, C-Y., Manson, J.E., Messina, C.R., Patel, M.J., Moreland, L., Stefanick, M.L., and LaCroix, A.Z. (2014). Sedentary Behaviour and Mortality in Older Women. *American Journal of Preventive Medicine*, **46(2)**, 122-135.

Sesso, H.D., Paffenbarger, R.S.Jr. and Lee, I.M. (2000). Physical activity and coronary heart disease in Men. *Circulation*, **102**, 975-980.

Shah, R.C., Buchman, A.S., Leurgans, S., Boyle, P.A. and Bennett, D.A. (2012). Association of total daily physical activity with disability in community-dwelling older persons: a prospective cohort study. *BMC Geriatrics*, **12**, 63.

Shepherd, R.J. (2001). Absolute versus relative intensity of physical activity in a dose-response context. *Medicine and Science in Sports and Exercise*, **33(6 Suppl)**, S400-S418.

Shepherd, R.J. and Fitcher, R. (1997). Physical activity and cancer: how may protection be maximised. *Critical Reviews in Oncology*, **8**, 219-272.

Shepherd, P.R. and Kahn, B.B. (1999). Glucose transporters and insulin action: implications for insulin resistance and diabetes mellitus. *New England Journal of Medicine*, **341**, 248-257.

Sigal, R.J., Kenny, G.P., Wasserman, D.H., Castaneda-Sceppa, C., and White, R.D. (2006). Physical activity/exercise and type 2 diabetes: a consensus statement from the American Diabetes Association. *Diabetes Care*, **29**, 1433-1438.

Siscovick, D.S., Fried, L., Mittelmark, M., Rutan, G., Bild, S. And O'Leary, D.H. (1997). Exercise intensity and subclinical cardiovascular disease in the elderly: the Cardiovascular Health Study. *American Journal of Epidemiology*, **145**, 977-986.

Skerrett, P.J. and Manson, J.E. (2002). Reduction in risk of coronary heart disease and diabetes. In *Handbook of Exercise in Diabetes*, edited by Ruderman, N., Devlin, J.T., Schneider, S.H., and Kriska, A. Alexandria, VA: American Diabetes Association, 2002.

Slattery, M.L., Edwards, S., Curtin, K., Ma, K., Edwards, R., Holubkov, R., and Schaffer, D. (2003). Physical activity and colorectal cancer. *American Journal of Epidemiology*, **158(3)**, 214-224.

Slattery, M.L. and Potter, J.D. (2002) Physical activity and colon cancer: confounding or interaction? *Medicine and Science in Sports and Exercise*, **34(6)**, 913-919.

Slattery, M.L., Edwards, S.L., Ma, K.N., Friedman, G.D., and Potter, J.D. (1997a). Physical activity and colon cancer: a public health perspective. *Annals of Epidemiology*, **7(2)**, 137-145.

Slattery, M.L., Potter, J., Caan, B., Edwards, S., Coates, A., Ma, K.N., and Berry, T.D. (1997b). Energy balance and colon cancer – beyond physical activity. *Cancer Research*, **57(1)**, 75-80.

Slattery, M.L., Schumacher, M.C., Smith, K.R., West, D.W., and Abd-Elghany, N. (1988). Physical activity, diet, and risk of colon cancer in Utah. *American Journal of Epidemiology*, **128**, 989-999.

Slentz, C.A., Houmard, J.A., Johnson, J.L., Bateman, L.A., Tanner, C.J., McCartney, J.S., Duscha, B.D., and Kraus, W.E. (2007). Inactivity, exercise training and detraining, and plasma lipoproteins.

STRIDE: a randomised, controlled study of exercise intensity and amount. *Journal of Applied Physiology*, **103**(2), 432-442.

Slentz, C.A., Aiken, L.B., Houmard, J.A., Bales, C.W., Johnson, J.L., Tanner, C.J., Duscha, B.D., and Kraus, W.E. (2005) Inactivity, exercise, and visceral fat. STRIDE: a randomised, controlled study of exercise intensity and amount. *Journal of Applied Physiology*, **99**, 1613-1618.

Smith, J.C., Nielson, K.A., Woodard, J.L., Seidenberg, M., Durgerian, S., Hazlett, K.E., Figueroa, C.M., Kandah, C.C., Kay, C.D., Matthews, M.A., and Rao, S.M. (2014). Physical activity reduces hippocampal atrophy in elders at genetic risk for Alzheimer's disease. *Frontiers in Aging Neuroscience*, **6**, doi: 10.3389/fnagi.2014.00061

Smith, P.J., Blumenthal, J.A., Hoffman, B.M., Cooper, H., Strauman, T.A., Welsh-Bohmer, K., Brown dyke, J.N., and Sherwood, A. (2010). Aerobic exercise and neurocognitive performance: a meta-analytic review of randomised controlled trials. *Psychosomatic Medicine*, **72**(3), 239-252.

Smith, R. (1985) Exercise and osteoporosis. *British Medical Journal*. **290**, 1163-1164.

Smith, S.M., Davis-Street, J.E., Fesperman, J.V., Calkins, D.S., Bawa, M., Macias, B.R., and Hargens, A.R. (2003). Evaluation of treadmill exercise in a lower body negative pressure chamber as a countermeasure for weightlessness-induced bone loss: a bed rest study with identical twins. *Journal of Bone Mineral Research*, **18**(12), 2223-2230.

Smorawiński, J., Kaciuba-Uściłko, H., Nazar, K., Kubala, P., Kamińska, E., Ziemia, A.W., Adrian, J., and Greenleaf, J.E. (2000). Effects of three-day bed rest on metabolic, hormonal and circulatory responses to an oral glucose load in endurance or strength trained athletes and untrained subjects. *Journal of Physiology and Pharmacology*, **51**(2), 279-289.

Smutok, M.A., Reece, C., Kokkinos, P.F., Farmer, C., Dawson, P., Shulman, R., DeVane-Bell, J., Patterson, J., Charabogios, C., Goldberg, A.P., and Hurley, B.F. (1993) Aerobic versus strength training for risk factor intervention in middle-aged men at high risk for coronary heart disease. *Metabolism*, **42**, 177-184.

Snow-Hunter, C., Whalen, R., Myburgh, K., Arnaud, S., and Marcus, R. (1992). Bone mineral density, muscle strength, and recreational exercise in men. *Journal of Bone and Mineral Research*, **7**, 1291-1296.

Sofi, F., Valecchi, D., Bacci, D., Abbate, R., Gensini, G.F., Casini, A., and Macchi, C. (2011). Physical activity and risk of cognitive decline: a meta-analysis of prospective studies. *Journal of Internal Medicine*, **269**, 107-117.

Sone, T., Miyake, M., Takeda, N., Tomomitsu, T., Otsuka, N., Fukunaga, M. (1996). Influence of exercise and degenerative vertebral changes on BMD: a cross-sectional study in Japanese men. *Gerontology*, **42**, 57-66.

Spate-Douglas, T. and Keyser, R.E. (1999). Exercise intensity: its effect on the HDL profile. *Archives of Physical Medicine and Rehabilitation*, **80**, 691-695.

Sport and Recreation Alliance (2012). Game of life: How sport and recreation can make us happier, healthier and richer.

http://www.sportandrecreation.org.uk/sites/sportandrecreation.org.uk/files/web/Game_of_Life/33_10_SRA_literary%20%review_v9%20WITH%20HYPERLINK.pdf

St Jeor, S.T., Brunner, R.L., Harrington, M.E. Scott, B.J., Daugherty, S.A., Cutter, G.R., Brownell, K.D., Dyer, A.R., and Foreyt, J.P. (1997). A classification system to evaluate weight maintainers, gainers, and losers. *Journal of the American Dietetic Association*, **97(5)**, 481-488.

Stephens, B.R., Grandados, K., Zderic, T.W., Hamilton, M.T., and Braun, B. (2011). Effect of 1 day of inactivity on insulin action in healthy men and women: interaction with energy intake. *Metabolism*, **60(7)**, 941-949.

Stevens, J., Truesdale, K.P., McCain, J.E., and Cai, J. (2006). The definition of weight maintenance. *International Journal of Obesity (London)*, **24(4)**, 395-403.

Stuart, C.A., Shangraw, R.E., Prince, M.J., Peters, E.J., and Wolfe, R.R. (1988). Bet-rest-induced insulin resistance occurs primarily in muscle. *Metabolism*, **37**, 802-806.

Standl, E. (2014). Statement accompanying World Diabetes Day, 2014.

Sumic, A., Michael, Y.L., Carlson, N.E., Howieson, D.B., and Kaye, J.A. (2007). Physical activity and the risk of dementia in oldest old. *Journal of Aging and Health*, **19(2)**, 242-259.

Sunami Y., Motoyama, M., Kinoshita, F., Mizooka, Y., Sueta, K., Matsunaga, A., Sasaki, J., Tanaka, H., Shindo, M. (1999). Effects of low-intensity aerobic training on the high-density lipoprotein cholesterol concentration in healthy elderly subjects. *Metabolism*, **48(8)**, 984-9.

Suominen, H. and Rahkila, P. (1991). Bone mineral density of the calcaneus in 70- to 80-yr-old male athletes and a population sample. *Medicine and Science in Sports and Exercise*, **23**, 1227-1233.

Swartz, A.M., Strath, S.J., Bassett, D.R., Moore, J.B., Redwine, B.A., Groer, M., and Thompson, D.L. (2003). Increasing daily walking improves glucose tolerance in overweight women. *Preventive Medicine*, **37**, 356-362.

Swedish National Institute of Public Health (2010). *Physical Activity in the Prevention and Treatment of Disease*. Professional Associations for Physical Activity (Sweden). Available at: <http://www.fyss.se/fyss-in-english/>

Szymanski, L.M. and Pate, R.R. (1994). Effects of exercise intensity, duration and time of day on fibrinolytic activity in physically active men. *Medicine and Science in Sports and Exercise*, **26**, 1102-1108.

Tanasescu, M., Leitzmann, M.F., Rimm, E.B., Willett, W.C., Stampfer, M.J., and Hu, F.B. (2002). Exercise type and intensity in relation to coronary heart disease in men. *Journal of the American Medical Association*, **288(16)**, 1994-2000.

- Tashiro, E., Miura, S., Koga, M., Sasaguri, M., Ideishi, M., Ikeda, M., Tanaka, H., Shindo, M., Arakawa, K. (1993). Crossover comparison between the depressor effects of low and high work-rate exercise in mild hypertension. *Clinical and Experimental Pharmacology and Physiology*, **20(11)**, 689-696.
- Terry, P., Baron, J.A., Weiderpass, E., Yuen, J., Lichtenstein, P., and Nyrén, O. (1999). Lifestyle and endometrial cancer risk: a cohort study from the Swedish Twin Registry. *International Journal of Cancer*, **82(1)**, 38-42.
- Thompson, D., Peacock, O., Western, M., and Batterham, A.M. (2015). Multidimensional physical activity: an opportunity not a problem. *Exercise and Sport Sciences Reviews*, **43(2)**, 67-74.
- Thompson, D. and Batterham, A.M. (2013). Towards integrated physical activity profiling. *PLoS ONE*, **8(2)**, e56427. Doi: 10.1371/journal.pone.0056427.
- Thompson, D., Batterham, A.M., Markovitch, D., Dixon, N.C., Lund, A.J.S., and Walhin, P. (2009). Confusion and conflict in assessing the physical activity status of middle-aged men. *PLoS ONE*, **4(2)**, e4337. Doi:10.1371/journal.pone.0004337.
- Thune, I. (2010). Cancer. In: *Physical Activity in the Prevention and Treatment of Disease*. Professionals Associations for Physical Activity, Sweden. Elanders: Sweden.
- Thune, I. and Furberg, A-S. (2001). Physical activity and cancer risk: dose-response and cancer, all sites and site-specific. *Medicine and Science in Sports and Exercise*, **33(6)**, 530-550.
- Thune, I. and Lund, E. (1996). Physical activity and risk of colorectal cancer in men and women. *British Journal of Cancer*, **70**, 57-62.
- Thune, I., Brenn, T., Lund, E., and Gaard, M. (1997). Physical activity and the risk of breast cancer. *New England Journal of Medicine*, **336**, 1269-1275.
- Tian, Q., Erickson, K.I., Simonsick, E.M., Aizenstein, H.J., Glynn, N.W., Boudreau, R.M., Newman, A.B., Kritchevsky, S.B., Yaffe, K., Harris, T.B., and Rosano, C. (2014). Physical activity predicts microstructural integrity in memory-related networks in very old adults. *Journal of Gerontology: Series A, Biological Sciences and Medical Sciences*, **69(10)**, 1284-1290.
- Tremblay, M.S., Colley, R.C., Saunders, T.J., Healy, G.N., and Owen, N. (2010). Physiological and health implications of a sedentary lifestyle. *Applied Physiology, Nutrition and Metabolism*, **35**, 725-740.
- Tsuzuku, S., Ikegami, Y. and Yabe, K. (1998). Effects of high-intensity resistance training on bone mineral density in young male powerlifters. *Calcified Tissue International*. **63**, 283-286.
- Tuomilehto, J., Lindstrom, J., Eriksson, J.G., Valle, T.T., Hamalainen, H., Ilanne-Parikka, P., Keinonen-Kiukaanniemi, S., Laakso, M., Louheranta, A., Rastas, M., Salminen, V., and Uusitupa, M. (2001). Prevention of type 2 diabetes mellitus by changes in lifestyle among subjects with impaired glucose tolerance. *New England Journal of Medicine*, **344**, 1343-1350.
- Ueji, M., Ueno, E., Osei-Hyiaman, D., Takahashi, H., and Kano, K. (1998). Physical activity and the risk of breast cancer: A case-control study of Japanese women. *Journal of Epidemiology*, **8(2)**, 116-122.

UKActive (2014), Turning the tide of inactivity.

http://ukactive.com/downloads/managed/Turning_the_tide_of_inactivity.pdf.

U.S. Department of Health and Human Services (2008). *Physical Activity Guidelines Advisory Committee Report*, Part G. 2008.

U.S. Department of Health and Human Services (2008). *Physical Activity Guidelines for Americans*. <http://www.health.gov/paguidelines/guidelines/default.aspx>

Uusi-Rasi, K., Sievanen, H. Vuori, I., Pasanen, M., Heinonen, A., and Oja, P. (1998). Associations of physical activity and calcium intake with bone mass and size in healthy women at different ages. *Journal of Bone and Mineral Research*, **13**, 133-142.

Vatten, L.J., Nilsen, T.I., Romundstad, P.R., Droyvold, W.B., and Holmen, J. (2006). Adiposity and physical activity as predictors of cardiovascular mortality. *European Journal of Cardiovascular Prevention and Rehabilitation*, **13(6)**, 909-915.

Van der Ploeg, H.P., Chey, T., Korda, R.J., Banks, E., and Bauman, A. (2012). Sitting time and all-cause mortality risk in 222,497 Australian adults. *Archives of Internal Medicine*, **172(6)**, 494-500.

Verghese, J., LeValley, A., Hall, C.B., Katz, M.J., Ambrose, A.F., and Lipton, R.B. (2006). Epidemiology of gait disorders in community-residing older adults. *Journal of the American Geriatrics Society*, **54**, 255-261.

Verloop, J., Rookaus, M.A., van der Koy, K., and Leuwen, F.E. (2000). Physical activity and breast cancer risk in women. *Journal of the National Cancer Institute*, **92**, 128-135.

Voss, M.W., Prakash, R.S., Erickson, K.I., Basak, C., Chaddock, L., Kim, J.S., Alves, H., Heo, S., Szabo, A.N., White, S.M., Wójcicki, T.R., Mailey, E.L., Gothe, N., Olson, E.A., McAuley, E., and Kramer, A.F. (2010). Plasticity of brain networks in a randomised intervention trial of exercise training in older adults. *Frontiers in Aging Neuroscience*, **2**, doi: 10.1097/PSY.0b013e3181d14633.

Voss, M.W., Vivar, C., Kramer, A.F., and van Praag, H. (2013) Bridging animal and human models of exercise-induced brain plasticity. *Trends in Cognitive Sciences*, **17**, 525-544.

Wang, J.S., Jen, C.J., Kung, H.C., Lin, L.J., Hsiue, T.R., and Chen, H.I. (1994). Different effects of strenuous exercise and moderate exercise on platelet function in men. *Circulation*, **90**, 2877-2885.

Wannamethee, S.G. and Shaper, A.G. (1992). Physical activity and stroke in British middle aged men. *British Medical Journal*, **304**, 597-601.

Wannamethee, S.G., Shaper, A.G., and Alberti, K.G. (2000). Physical activity, metabolic factors, and the incidence of coronary heart disease and type 2 diabetes. *Archives of Internal Medicine*, **160**, 2108-2116.

Warburton, D.E., Charlesworth, S., Ivey, A., Nettlefold, L., and Bredin, S.S. (2010). A systematic review of the evidence for Canada's Physical Activity Guidelines for Adults. *International Journal of Behavioural Nutrition and Physical Activity*, **7**, 39.

- Wei, M., Gibbons, L.W., Mitchell, T.L., Kampert, J.B., Lee, C.D., and Blair, S.N. (1999). The association between cardiorespiratory fitness and impaired glucose and type 2 diabetes mellitus in men. *Annals of Internal Medicine*, **130(2)**, 89-96.
- Weinstein, A.M., Voss, M.W., Prakash, R.S., Chaddock, L., Szabo, A., White, S.M., Wojcicki, T.R., Mailey, E., McAuley, E., Kramer, A.F., and Erickson, K.I. (2012). The association between aerobic fitness and executive function is mediated by prefrontal cortex volume. *Brain, Behavior, and Immunity*, **26(5)**, 811-819.
- Weinstein, A.R., Sesso, H.D., Lee, I-M., Cook, N.R., Manson, J.E. Buring, J.E., and Gaziano, J.M. (2004). Relationship of physical activity vs body mass index with type 2 diabetes in women. *Journal of the American Medical Association*, **292**, 1188-1194.
- Weiss, C., Seitel, G. and Bartsch, P. (1998). Coagulation and fibrinolysis after moderate and very heavy exercise in healthy male subjects. *Medicine and Science in Sports and Exercise*, **30**, 246-251.
- Wen, C.P., Wai, J.P.M., Tsai, M.K., Yang, Y.C., Cheng, T.Y.D., Lee, M-C, Chan, H.T., Tsao, C.K., Tsai, S.P., and Wu, X.F. (2011) Minimum amount of physical activity for reduced mortality and extended life expectancy: a prospective cohort study. *Lancet*, **378(9798)**, 1244-1253.
- Wendel-Voss, G.C.W., Schuit, A.J., Feskens, E.J.M., Boshuizen, H.C., Verschuren, W.M.M., Saris, W.H.M., and Kromhout, D. (2004). Physical activity and stroke. A meta-analysis of observational data. *International Journal of Epidemiology*, **33(4)**, 787-798.
- Whelton, S.P., Chin, A., Xin, X., and He, J. (2002). Effect of aerobic exercise on blood pressure: a meta-analysis of randomised, controlled trials. *Annals of Internal Medicine*, **136(7)**, 493-503.
- Williams P.T. (2001). Physical fitness and activity as separate heart disease risk factors: a meta-analysis. *Medicine and Science in Sports and Exercise*, **33(5)**, 754-761.
- Williams P.T. (1998). Relationships of heart disease risk factors to exercise quantity and intensity. *Archives of Internal Medicine*, **158(3)**, 237-245.
- Williams P.T. (1996). High-density lipoprotein cholesterol and other risk factors for coronary heart disease in female runners. *New England Journal of Medicine*, **334(20)**, 1298-1303.
- Williams P.T., Wood, P.D., Haskell, W.L., Vranizan, K. (1982). The effects of running mileage and duration on plasma lipoprotein levels. *Journal of the American Medical Association*, **247(19)**, 2674-9.
- Wilmot, E.G., Edwardson, C.L., Achana, F.A., Davies, M.J., Gorely, T., Gray, L.J., Khunti, K., Yates, T., and Biddle, S.J.H. (2012) Sedentary time in adults and the association with diabetes, cardiovascular disease and death: a systematic review and meta-analysis. *Diabetologica*, **55**, 2895-2905.
- Wittich, A., Mautalen, C.A., Oliveri, M.B., Bagur, A., Somoza, F., and Rotemberg, E. (1998). Professional football (soccer) players have a markedly greater skeletal mineral content, density and size than age- and BMI-matched controls. *Calcified Tissues International*, **63**, 112-117.

Wood P.D., Haskell, W.L., Blair, S.N., Williams, P.T., Krauss, R.M., Lindgren, F.T., Albers, J.J., Ho, P.H., and Farguhar, J.W. (1983). Increased exercise level and plasma lipoprotein concentrations: a one year randomized, controlled study in sedentary middle-aged men. *Metabolism*, **32(1)**, 31-39.

Woodcock, J., Franco, O.H., Orsini, N., and Roberts, I. (2011). Non-vigorous physical activity and all-cause mortality: Systematic review and meta-analysis of cohort studies. *International Journal of Epidemiology*, **40**, 121-138.

World Cancer Research Fund, American Institute for Cancer Research (2007). *Food, nutrition, physical activity, and the prevention of cancer. A global perspective*. AICR, Washington DC.

World Cancer Research Fund/American Institute for Cancer Research (1997). *Food, nutrition and the prevention of cancer: a global perspective*. Washington, DC, American Institute for Cancer Research.

World Health Organisation (2009) *Global Health Risks: Mortality and burden of disease attributable to selected major risks*, Geneva, Switzerland: World Health Organisation.

World Health Organisation (2000). *Obesity: preventing and managing the global epidemic*. Geneva, Switzerland: IOTF, WHO Technical Report Series 894.

Yanagibori, R., Kondo, K., Suzuki, Y., Kawakubo, K., Iwamoto, T., Itamoto, H., and Gunji, A. (1998). Effect of 20 days' bed rest on the reverse cholesterol transport system in healthy young subjects. *Journal of Internal Medicine*, **243(4)**, 307-312.

Young, J.C., Enslin, J. and Kuca, B. (1989). Exercise intensity and glucose tolerance in trained and non-trained subjects. *Journal Applied Physiology*, **67**, 39-43.

Zhang, Y., Cantor, K.P., Dosemeci, M., Lynch, C.F., Zhu, Y., and Zheng, T. (2006). Occupational and leisure-time physical activity and risk of cancers of the colon and rectum: an Italian case-control study. *British Journal of Cancer*, **79(11-12)**, 1912-1916.

Zwart, S.R., Hargens, A.R., Lee, S.M., Macias, B.R., Watenpaugh, D.E., Tse, K., and Smith, S.M. (2007). Lower body negative pressure treadmill exercise as a countermeasure for bed rest-induced bone loss in female identical twins. *Bone*, **40(2)**, 529-537.